

10049957 021502
JC07 Rec'd PCT/PTO 19 FEB 2002

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER KATO=21
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. APPLICATION NO. (If known, see 37 CFR 1.5) 10/049957
INTERNATIONAL APPLICATION NO. PCT/JP00/05590	INTERNATIONAL FILING DATE 21 August 2000	PRIORITY CLAIMED 19 August 1999
TITLE OF INVENTION CHONDROGENESIS PROMOTERS		
APPLICANT(S) FOR DO/EO/US Yukio KATO et al.		
<p>Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information.</p> <ol style="list-style-type: none"> <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371 <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). <input checked="" type="checkbox"/> The US has been elected in a Demand by the expiration of 19 months from the priority date (PCT Article 31). <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> <input type="checkbox"/> is attached hereto (required only if not transmitted by the International Bureau) <input checked="" type="checkbox"/> has been communicated by the International Bureau. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US) <input checked="" type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)) <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) <ol style="list-style-type: none"> <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> have been communicated by the International Bureau. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. <input checked="" type="checkbox"/> have not been made and will not be made. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)) <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)) <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). <p>Items 11. to 16. below concern document(s) or information included:</p> <ol style="list-style-type: none"> <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98 <input type="checkbox"/> An Assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. <input type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. <input type="checkbox"/> A substitute specification <input type="checkbox"/> A change of power of attorney and/or address letter <input checked="" type="checkbox"/> Other items or information <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Courtesy copy of the first page of the International Publication (WO 01/13951) <input checked="" type="checkbox"/> Courtesy copy of the International Preliminary Examination Report (In Japanese). <input checked="" type="checkbox"/> Formal drawings, 6 sheets, Figures 1-7 <input checked="" type="checkbox"/> Courtesy Copy of the International Search Report <p><input checked="" type="checkbox"/> The application is (or will be) assigned to CHUGAI SEIYAKU KABUSHIKI KAISHA, whose address is 5-1, Ukima 5-chome, Kita-ku, Tokyo 115-8543, Japan</p>		

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)
10/049957International Application No
PCT/JP00/05590Attorney's Docket No
KATO=21

17. [xx] The following fees are submitted

BASIC NATIONAL FEE (37 CFR 1.492 (a)(1)-(5):Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO. **\$1040.00**International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO .. **\$890.00**International preliminary examination fee (37 CFR 1.482) not paid to USPTO but
international search fee (37 CFR 1.445(a)(2)) paid to USPTO. **\$740.00**International preliminary examination fee paid to USPTO (37 CFR 1.482)
but all claims did not satisfy provisions of PCT Article 33(1)-(4) **\$710.00**International preliminary examination fee paid to USPTO (37 CFR 1.482)
and all claims satisfied provisions of PCT Article 33(1)-(4)... .. **\$100.00****ENTER APPROPRIATE BASIC FEE AMOUNT =****CALCULATIONS PTO USE ONLY**

\$ 890.00

Surcharge of **\$130.00** for furnishing the oath or declaration later than [] 20 [] 30
months from the earliest claimed priority date (37 CFR 1.492(e)).

Claims as Originally Presented

Number Filed

Number Extra

Rate

Total Claims 24 - 20 04 X \$18 00

Independent Claims 6 - 3 03 X \$84 00

Multiple Dependent Claims (if applicable) +\$280.00

TOTAL OF ABOVE CALCULATIONS =

Claims After Post Filing Prel Amend

Number Filed

Number Extra

Rate

Total Claims - 20 X \$18.00

Independent Claims - 3 X \$84 00

TOTAL OF ABOVE CALCULATIONS =Reduction of 1/2 for filing by small entity, if applicable. Applicant claims small entity
status. See 37 CFR 1.27**SUBTOTAL =**Processing fee of **\$130.00** for furnishing the English translation later than [] 20 [] 30
months from the earliest claimed priority date (37 CFR 1.492(f)).**TOTAL NATIONAL FEE =**Fee for recording the enclosed assignment (37 CFR 1.21(h)) The assignment must be
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) **\$40.00** per property +**TOTAL FEES ENCLOSED =**Amount to be:
refunded \$
charged \$

- a. [] A check in the amount of \$_____ to cover the above fees is enclosed
- b. [X] Credit Card Payment Form (PTO-2038), authorizing payment in the amount of \$ 1,494 00, is attached.
- c. [] Please charge my Deposit Account No. **02-4035** in the amount of \$_____ to cover the above fees.
A duplicate copy of this sheet is enclosed
- d. [XX] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment
to Deposit Account No. **02-4035** A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.437(a) or
(b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO.

BROWDY AND NEIMARK, P.L.L.C.
624 NINTH STREET, N.W., SUITE 300
WASHINGTON, D.C. 20001

TEL: (202) 628-5197

FAX: (202) 737-3528

Date of this submission: **February 19, 2002**

Form PTO-1390 (as slightly revised by Browdy and Neimark)

SIGNATURE

Sheridan Neimark

NAME

20,520

REGISTRATION NUMBER

6/p/s

SPECIFICATION

CHONDROGENESIS STIMULATORTECHNICAL FIELD

This invention relates to a novel chondrogenesis
5 stimulator. More specifically, the invention relates to a
chondrogenesis stimulator containing a membrane-bound
transferrin-like protein (hereunder sometimes referred to
as MTf).

PRIOR ART

10 The cartilage tissue of animals is composed of
chondrocytes and matrix. The cartilage tissue accounts
for the greater part of the skeleton at the prenatal stage
and it is postnatally replaced with bone tissue due to
endochondral ossification. When endochondral ossification
15 starts, chondrocytes change from the resting to
proliferating phase and the proliferating chondrocytes
are then differentiated into hypertrophic chondrocytes
[Reference;, "Hone no kagaku (Science of Bone)", ed. by
Tsuda et al., pp. 11-29, Tokyo Ishiyaku Shuppan, 1982].
20 Thus, it has been well known that chondrocytes are
essential cells for the formation of bone tissue,
particularly at the growth stage. However, the
differentiation of chondrocytes and the endochondral
ossification remain unknown in many aspects.

25 The cell membrane of chondrocytes has characteristic
glycoproteins and their membrane proteins might contribute
to the unique features of chondrocytes that distinguish
them from the cells of other connective tissues (as

exemplified by spherical cell morphology, massive secretion of cartilage matrix, survival and proliferation in soft agar). Based on this hypothesis, Yan et al. (Yan et al.; J. Biol. Chem., vol. 265, pp. 10125-10131, 1990) and Kato et al. (Kato et al., Journal of the Society of Bone Metabolism of Japan, vol. 10, No. 2, pp. 187-192, 1992) investigated the effects of various lectins on the differentiation and proliferation of chondrocytes and, among other things, they have shown that concanavalin A (hereunder sometimes referred to as Con A) which is Jack bean lectin and which has affinity for α -D-mannose residue and α -D-glucose residue is a potent stimulator of chondrogenic differentiation, with the increase in proteoglycan synthesis being one of the criteria for the Con A activity. Chondrocytes treated with Con A change their shape from the immature flat morphology to the differentiated spherical form, inducing the production of proteoglycan and type II collagen which are markers of chondrogenic differentiation, the expression of alkaline phosphatase, etc., and even the calcification. Other lectins do not exert such differentiation inducing action.

In an attempt to search for a receptor mediating the action of Con A, Kawamoto et al. (Kawamoto et al., Eur. J. Biochem. vol. 256, pp. 503-509, 1998) paid particular attention to a protein of 76 kDa (p76) which was one of the about 20 kinds of Con A-binding proteins on chondrocytes and which would be expressed at lower levels in retinoic acid treated chondrocytes (upon treatment with retinoic

acid, chondrocytes are dedifferentiated to lose reactivity with Con A). After purifying p76 from the plasma membrane fraction of rabbit chondrocytes by Con A affinity column chromatography, the N-terminal amino acid sequence was
5 determined and the gene was cloned. In view of the determined amino acid sequence and the nucleic acid sequence of its cDNA, p76 showed 86% amino acid identity with melanotransferrin (p97) and was considered its counterpart; p97 is known as a tumor-associated antigen
10 expressed at high levels in human tumors such as melanoma. The physiological functions of p97 are yet to be known and its expression has been reported to be high in only tumor cells, with very low detectability in normal tissue.

In view of its ability to bind with Con A, p76 is
15 presumably involved in the differentiation of chondrocytes or in the development of their function; however, nothing has been confirmed about the effects this protein would actually impose on chondrocytes or their precursors.

An object, therefore, of the present invention is to
20 identify a substance that will be involved in the differentiation of chondrocytes and provide a novel chondrogenesis stimulator using the substance. The present invention will lead to the invention of a substance that can control the function of chondrocytes and which
25 eventually enables promoted osteogenesis. The substance can potentially lead to the treatment, prevention and diagnosis of new types of diseases associated with the cartilage and bone metabolisms.

DISCLOSURE OF THE INVENTION

In order to attain the stated object, the present inventors made intensive studies and found that differentiation to cartilage could be markedly induced by overexpressing a membrane-bound transferrin-like protein (MTf) gene in mouse cell line ATDC5 which retained the ability to differentiate to chondrocytes but which would hardly differentiate in the absence of stimulation.

Thus, the present invention provides a chondrogenesis stimulator containing a membrane-bound transferrin-like protein (MTf).

The MTf is preferably rabbit p76 protein, human p97 protein, mouse MTf protein, as well as a protein demonstrating the MTf activity that has an amino acid sequence encoded by DNA which hybridizes, under stringent conditions, with a DNA that encodes p76 protein or p97 protein or mouse MTf, and human p97 protein is particularly preferred.

The MTf is most preferably selected from the following:

- 1) a protein having the amino acid sequence of SEQ ID NO: 2;
- 2) a protein having the amino acid sequence of SEQ ID NO: 4;
- 3) a protein having the amino acid sequence of SEQ ID NO: 15; and
- 4) a protein demonstrating the MTf activity that has an amino acid sequence encoded by DNA which hybridizes, under

stringent conditions, with a DNA encoding the protein of
SEQ ID NO: 2, 4 or 15.

The present invention also provides said
chondrogenesis stimulator in which the MTf lacks the GPI
5 anchor region.

The chondrogenesis stimulator of the invention
becomes more effective when used in combination with an MTf
activating agent and/or insulin.

The chondrogenesis stimulator of the invention is
10 useful with the following diseases: OA (osteoarthritis);
RA (rheumatoid arthritis); injury of articular cartilage
due to trauma; maintenance of chondrocyte phenotype in
autologous transplantation of chondrocytes; reconstruction
of cartilage in the ear, trachea or nose; osteochondritis
15 dissecans; regeneration of intervertebral disk or meniscus;
fractured bone; and osteogenesis from cartilage.

The invention further provides an agent for gene
therapy to promote chondrogenesis which contains as an
active ingredient an expression vector incorporating a DNA
20 coding for any one of the following proteins:

- 1) a protein having the amino acid sequence of SEQ ID NO:
2;
- 2) a protein having the amino acid sequence of SEQ ID NO:
4;
- 25 3) a protein having the amino acid sequence of SEQ ID NO:
15; 4) a protein demonstrating the MTf activity that has an
amino acid sequence encoded by DNA which hybridizes, under
stringent conditions, with a DNA encoding the protein of

SEQ ID NO: 2, 4 or 15; and

5) a protein which is the same as protein 1), 2), 3) or 4), except that it lacks the GPI anchor region.

The present invention further provides a chondrogenic
5 differentiation suppressing agent containing an MTf antagonist.

The MTf antagonist is preferably an anti-MTf antibody or an oligonucleotide or an oligonucleotide analog that are hybridizable with a nucleic acid encoding MTf.

10 The present invention further provides a method for screening an MTf activating agent which comprises the steps of:

1) preparing a cell line in which MTf is overexpressed, wherein said cell line retains the ability to differentiate
15 to chondrocytes but hardly differentiate without stimulation;

2) adding candidate substances to the cell line prepared in step 1) and culturing it for a specified period of time; and

20 3) examining the cell line for induced chondrogenic differentiation and selecting an MTf activating agent from the candidate substances.

The present invention also provides an MTf activating agent as obtained by the method described above.

25 The present invention also provides a chondrogenesis stimulator containing an MTf activating agent as obtained by the method described above.

The present invention further provides MTf which

lacks the GPI anchor region.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows a scheme of the procedure of preparing an MTf overexpressing ATDC5 variant cells;

5 Fig. 2 shows the expression of the MTf gene in the variant of ATDC5 cells as analyzed by Northern blotting (photographs of electrophoresis);

Fig. 3 shows the expression of the MTf protein in the variant of ATDC5 cells as analyzed by Western blotting
10 (photographs of electrophoresis);

Fig. 4 is a set of photographs showing that MTf overexpressing cell lines (Full-1 and Full-5) demonstrate the morphology of differentiated chondrocytes in comparison with control cells (pC-1), all of which are cultured for 29
15 days in the absence of insulin (biological morphology is shown);

Fig. 5 is a set of photographs showing that MTf overexpressing cell lines (Full-1 and Full-5) demonstrate the morphology of differentiated chondrocytes in comparison
20 with control cells (pC-1), all of which are cultured for 29 days in the presence of insulin (biological morphology is shown);

Fig. 6 is a set of photographs showing the effects of the addition of the conditioned medium of rabbit
25 chondrocytes on the induction of chondrogenic differentiation (biological morphology is shown); and

Fig. 7 shows the result of RT-PCR Southern blotting which demonstrates the overexpression of antisense MTf RNA

and the suppression of aggrecan synthesis in the presence and absence of insulin.

BEST MODE FOR CARRYING OUT THE INVENTION

The term "membrane-bound transferrin-like protein
5 (MTf)" as used in the invention means a protein on the cell membrane of chondrocytes that binds to Con A and which has iron-binding sites as transferrin does. Preferably, the term means a protein having the ability to mediate the induction of chondrogenic differentiation by Con A.

10 The term MTf has conventionally been used as the abbreviation for melanotransferrin (p97) known as a tumor antigen expressed at high levels in melanoma and other tumors. As it turned out, however, p97 is also expressed at high levels in tissues other than cancer, particularly
15 in cartilage. Since p97 is by no means specific to cancer tissue, the present inventors redefined the term MTf as meaning "membrane-bound transferrin-like protein".

The term "MTf activity" as used herein means an activity that induces undifferentiated cells to
20 differentiate to cartilage and which promotes chondrocytes to develop their function.

Examples of MTf include but are not limited to rabbit p76 protein, p97 protein which is a human protein homologous to rabbit p76 protein, mouse MTf protein, as
25 well as proteins having MTf activity that contain alterations such as deletion, substitution or addition of one or more of the amino acids of these proteins, and proteins having MTf activity amino acid sequences encoded

by DNA which hybridizes with DNA encoding p76 protein or p97 protein or mouse MTF protein under stringent conditions [a typical method is described in Molecular Cloning: A Laboratory Manual, Sambrook et al., Cold Spring Harbor Laboratory Press, 1989 and consists, for example, of hybridization at 68°C in a solution containing 6 x SSC, 0.5% SDS, 10 mM EDTA, 5 x Denhardt's solution and 10 mg/ml of denatured salmon sperm DNA].

Rabbit p76 protein is homologous to human p97 protein and sometimes called rabbit p97 (Kawamoto et al., Eur. J. Biochem. vol. 256, pp. 503-509, 1998). The nucleotide and amino acid sequences of rabbit p76 protein are identified by SEQ ID NO: 1 and SEQ ID NO: 2, respectively. The nucleotide and amino acid sequences of human p97 protein are also known (Rose, T.M. et al., Proc. Natl. Acad. Sci. USA 83, 1261-1265, 1986). The nucleotide and amino acid sequences of human p97 protein are identified by SEQ ID NO: 3 and SEQ ID NO: 4, respectively. Mouse MTF protein is described in Biochim. Biophys. Acta, 1447:258-264, 1999 and its nucleotide and amino acid sequences are identified by SEQ ID NO: 14 and SEQ ID NO: 15, respectively. The homology between the MTF proteins over animal species are high and the amino acid identity is 83% between mouse and human, 82% between mouse and rabbit, and 86% between human and rabbit.

p76/p97 proteins are GPI anchored proteins which have glycolipid GPI (glycosylphosphatidylinositol) bound to the carboxyl group in C-terminal amino acid so that they are bound to membranes using GPI as an anchor (for p76, see Ryo

Oda, Journal of Dentistry, Hiroshima University, vol. 29, No. 1, pp. 40-57, 1997; for p97, see Alemany, R. et al., J. Cell Science, 104, 1155-1162, 1993). As will be shown later in the Examples, it was verified that not only
5 full-length MTf but also GPI anchor lacking MTf or soluble MTf have a chondrogenic differentiation inducing activity when they were expressed in non-MTf-expressing cells. Therefore, such soluble MTf, preferably the GPI anchor lacking MTf, can also be used as a chondrogenesis
10 regulating agent. The GPI anchor lacking MTf as used herein means a soluble MTf which lacks all or part of the GPI anchor moiety; in the case of rabbit MTf, it may be exemplified by MTf in which the 28 residues at C-terminal necessary for GPI anchor binding are deleted and in the
15 case of human MTf and mouse MTf, it may be exemplified by MTf in which the 30 residues at C-terminal necessary for GPI anchor binding are deleted.

The MTf to be used in the invention may be of a native or recombinant form and either form can be obtained
20 by methods known in the art. The respective types of MTf are illustrated below.

Native form

MTf can be prepared by the method described in JP 7-82297A using chondrocytes. Briefly, cartilage tissues of
25 various animals can be used as chondrocyte source; for example, a rabbit costal growth plate cartilage as the source is treated with protease and collagenase in accordance with the method of Kato et al. (Kato et al.; J.

Cell Biol., vol. 100, pp. 477-485, 1985) to obtain chondrocytes. The isolated chondrocytes can be incubated in a medium containing fetal calf serum (FCS) on a culture dish at 37°C in an atmosphere of 5% CO₂ and 95% air. The
5 cultured chondrocytes are recovered, homogenized with a homogenizer and subjected to sedimentation equilibrium centrifugation by 17%/40% sucrose equilibrium density gradient to separate membrane proteins. The obtained membrane protein fraction is directly subjected on a
10 concanavalin A affinity column; alternatively, in order to remove membrane proteins that bind to lectins other than concanavalin A, the membrane protein fraction is first subjected on an affinity column of wheat germ lectin which is a typical lectin and then subjected on a concanavalin A
15 affinity column. By these and other techniques, more of the concanavalin A binding protein fraction can be separated. The specificity of the obtained concanavalin A bound protein fractions for chondrocytes can be evaluated by comparing these fractions through SDS-polyacrylamide gel
20 electrophoresis (SDS-PAGE). After identifying the desired chondrocyte specific glycoproteins, bands of interest are excised from the gel, extracted and purified by electroelution or other suitable techniques. The resulting glycoproteins can be analyzed for the sugar chains after
25 excising by endoglicosidase.

Recombinant form

Recombinant MTf can be prepared by the methods described in the Examples of the invention or modifications

thereof; by these methods, plasmids incorporating the MTf gene are transfected to host cells for expressing the MTf protein.

5 However, these are not the only methods that can be used and various methods of transformation and various host cells that are known in the art can also be used. For example, a gene encoding MTf may be inserted into a suitable vector to transform prokaryotic or eukaryotic host cells.

10 Further, suitable stimulators or sequences that are involved gene expression may be introduced into the vectors to enable gene expression in the transformed host cells. If desired, a gene of interest may be linked to genes encoding other polypeptides to express it as a fused
15 protein so that it can be purified with greater ease or expressed at higher level. The desired protein can also be excised by applying suitable treatments in the purification step.

It is generally held that eukaryotic genes show
20 polymorphism as is known for the human interferon gene. The polymorphism may cause substitution by one or more amino acids or it may cause changes in base sequences with no change occurring in amino acids.

Even polypeptides having deletion or addition of one
25 or more amino acids within the amino acid sequence of SEQ ID NO: 2, 4 or 15 or having substitution of one or more amino acids may have a cell cycle regulating activity. For example, it is already known that a polypeptide having

substitution of cysteine for serine in the human interleukin 2 (IL-2) which is derived from nucleotide alterations has exerted an IL-2 activity (Wang et al., Science 224:1431, 1984). These techniques for preparing
5 modified genes encoding MTf protein are known to the skilled artisan.

In many cases of expression in eukaryotic cells, sugar chains may be added to the protein and the addition of sugar chains can be regulated by substituting one or
10 more amino acids of the protein and even in this case, the chondrogenic differentiation inducing activity may be exhibited. Therefore, genes encoding such polypeptides obtained by using artificial modifications of the gene encoding MTf gene can all be used in the invention, as long
15 as such polypeptides have the chondrogenic differentiation inducing activity.

Expression vectors that can be used include replication origins, selection markers, promoters, RNA splicing sites, polyadenylation signals and so on.

20 Prokaryotic organisms that can be used as host cells in the expressing system include, for example, *Escherichia coli* and *Bacillus subtilis*. Eukaryotic microorganisms that can be used as host cells include, for example, yeasts and myxomycetes. If desired, insect cells such as Sf9 may be
25 used as host cells. Host cells derived from animal cells include, for example, COS cells and CHO cells.

Transformants thus obtained by transforming with the gene encoding MTf protein are cultured to produce proteins.

insulin. However, it was found that the effect of the chondrogenesis stimulator is further enhanced in the presence of insulin. Therefore, the desired cartilage repairing action could be further enhanced by using MTf in
5 combination with MTf activating agents such as insulin and an insulin-like growth factor.

When the supernatant of a chondrocyte culture was added, marked differentiation of chondrocytes was observed in MTf overexpressing cell lines, suggesting that an MTf
10 activating agent may exist in the conditioned medium of a chondrocyte culture. Therefore, the desired cartilage repairing action could be further potentiated by using MTf in combination with an MTf activating agent.

The MTf activating agent may be obtained by the
15 following methods:

- 1) purifying from the conditioned medium of a chondrocyte culture system;
- 2) cloning the cDNA for protein binding to an MTf from a chondrocyte cDNA library; and
- 20 3) cloning the cDNA of protein binding to an MTf by the yeast two-hybrid method.

To screen various candidate substances for an MTf activating agent, a method including the following steps can be used:

- 25 1) preparing a cell line in which MTf is overexpressed, wherein said cell line retains the ability to differentiate to chondrocytes but hardly differentiate without stimulation;

MTf protein or an MTf mutant (variant) introduced in it.

The agent for gene therapy of the invention contains as an active ingredient an expression vector incorporating a DNA coding for any one of the following proteins:

- 5 1) a protein having the amino acid sequence of SEQ ID NO: 2;
- 2) a protein having the amino acid sequence of SEQ ID NO: 4;
- 3) a protein having the amino acid sequence of SEQ ID NO: 15;
- 10 4) a protein demonstrating the MTf activity that has an amino acid sequence encoded by DNA which hybridizes, under stringent conditions, with a DNA encoding the protein of SEQ ID NO: 2, 4 or 15; and
- 5) a protein which is the same as protein 1), 2), 3) or 4),
15 except that it lacks the GPI anchor region.

DNAs encoding MTf variants can be easily prepared by the skilled artisan using known techniques such as site-directed mutagenesis and PCR [Molecular Cloning: A Laboratory Manual, 2nd ed., Chapter 15, Cold Spring Harbor
20 Laboratory Press (1989), and PCR - A Practical Approach, IRL Press, 200-210 (1991)].

In the present invention, an MTf- or MTf variant expression vector is provided as a DNA to be introduced into cells. These expression vectors can be prepared by
25 linking DNA encoding an MTf- or MTf variant to an expression vector such as pSG5 (Stratagene). In the next step, the prepared DNA mixture is introduced into cells. Exemplary cells may include bone marrow interstitial cells,

fibroblasts, periosteal cells, perichondral cells, synovial cells and dedifferentiated chondrocytes. DNA can be introduced into cells by the calcium phosphate method [Idenshi donyu to hatsugen kaisekiho (Gene Introduction and Methods of Expression and Analysis), ed. by Takashi Yokota and Kenichi Arai, Yodosha, 1994]. Hence, by using the introduced DNA as a medicinal active ingredient, one can prepare an agent for gene therapy which has the chondrogenesis promoting action. It is thought that, upon administering such agent for gene therapy, MTF or its variant would be expressed at high levels in cells, promoting the action of inducing chondrogenic differentiation in the cells. Therefore, the MTF containing agent of the invention for gene therapy can be used as a therapeutic or preventive of the various diseases listed above.

The agent of the invention for gene therapy can be introduced into cells by either a virus vector based method of gene introduction or a non-viral method of gene introduction [Nikkei Science, April 1994, pp. 20-45, Jikken igaku zokan (Extra Issue of Experimental Medicine), 12(15)(1994), and Jikken igaku bessatsu (Supplement to Experimental Medicine), "Idenshi chiryo no kiso gijutsu (Basic Technology in Gene Therapy)", Yodosha (1996)].

In an example of the viral vector based method of gene introduction, DNA encoding an MTF or a variant MTF is inserted into DNA or RNA viruses such as retrovirus, adenovirus, adeno-associated virus, herpes virus, vaccinia

virus, poxvirus, poliovirus and Sindbis virus. Non-viral methods of gene introduction include direct intramuscular administration of an expression plasmid (DNA vaccination), liposome method, lipofectin method, microinjection, calcium phosphate method, and electroporation.

In order to ensure that the agent for gene therapy of the invention acts as a practical medicine, two methods may be used, that is, an in vivo method where DNA is directly introduced into the body and an ex vivo method where a certain kind of cells are taken out of a human and DNA is introduced into the cell, which is then put back into the body [Nikkei Science, April 1994, pp. 20-45, Gekkan yakuji (Monthly Yakuji), 36(1), 23-48 (1994), and Jikken igaku zokan (Extra Issue of Experimental Medicine), 12(15)(1994)]. The in vivo method is more preferred.

When administering the agent for gene therapy by the in vivo method, the route of administration should depend on the disease, its severity and other factors. Exemplary methods of administration include intra-articular injection, direct application to a missing part of articular cartilage, implantation (with putty, polylactic acid, etc.) and intra-articular sustained-release agent. Intravenous injection is also possible. For administration by the in vivo method, injections are generally used, with conventional carriers being added as required. Liposomes or membrane fused liposomes may be formulated as suspensions, frozen vesicles, centrifugally concentrated frozen vesicles.

amino acid sequence identified by SEQ ID NO: 2, 4 or 15.

The term "oligonucleotide" as used herein means oligonucleotides generated from naturally occurring bases and sugar portions bound by intrinsic phosphodiester bonds, as well as analogs thereof. Therefore, the first group encompassed by this term comprises naturally occurring species or synthetic species that are generated from naturally occurring subunits or homologs thereof. The term "subunit" means a base-sugar combination which links to adjacent subunit by phosphodiester bond or other bond. The second group of oligonucleotides are their analogs that function similar to oligonucleotides but which are composed of residues having non-naturally-occurring moieties. These include oligonucleotides having chemical modifications applied to phosphate groups, sugar portions and 3'- and 5'-ends in order to provide increased stability. Examples are oligophosphorothioate and oligomethylphosphonate in which one of the oxygen atoms in the phosphodiester group between nucleotides is substituted by sulfur and $-CH_3$, respectively. Phosphodiester bonds may be replaced by other structures which are non-ionic and achiral. Additional oligonucleotide analogs that can be used are species containing modified base forms, that is, purine and pyrimidine in non-naturally-occurring form.

The oligonucleotides to be used in the invention have preferably 5 - 40 subunits in length, more preferably 8 - 30 subunits, most preferably 12 - 30 subunits.

In the present invention, the target portion of mRNA

with which oligonucleotides hybridize is preferably a transcription initiation site, a translation initiation site, an intron/exon junction site or a 5'-cap site; considering the secondary structure of mRNA, a site having
5 no steric hindrance should be selected.

In the present invention, peptide nucleic acids (see, for example, Bioconjugate Chem., Vol. 5, No. 1, 1994) may be used in place of oligonucleotides.

In a particularly preferred embodiment of the
10 invention, oligonucleotides or peptide nucleic acids that hybridize with a nucleotide sequence encoding the amino acid sequence identified by SEQ ID NO: 2 and which can inhibit MTF expression is employed.

In the present invention, oligonucleotides can be
15 produced by synthesis methods known in the art, for example, the solid-phase synthesis method using a synthesizer as manufactured by Applied Biosystems. Similar methods can be used to produce oligonucleotide analogs such as phosphorothioate and alkylated derivatives [Akira
20 Murakami et al., "Kinosei antisense DNA no gosei (Chemical Synthesis of Functional Antisense DNA)", Organic Synthesis Chemistry, 48(3):180-193, 1990].

The MTF antagonist that can be used in the invention is not limited to oligonucleotides of the above-defined
25 antisense DNA providing length. To the extent that production of intrinsic MTF can be suppressed, a longer antisense, preferably an antisense of 500 - 600 nucleotides in length, may be inserted into a genome to be used for

suppressing chondrogenic differentiation (see Example 3).

The anti-MTf antibody to be used in the invention is one that recognizes a peptide having at least five consecutive amino acids in the amino acid sequence identified by SEQ ID NO: 2, 4 or 15; this can be produced using a conventional procedure [see, for example, Shin-seikagaku jikken koza 1 (New Course in Biochemical Experiments 1), Protein I, pp. 389-397, 1992], which comprises immunizing an animal with an antigenic peptide having at least five consecutive amino acids in the amino acid sequence of SEQ ID NO: 2, 4 or 15, isolating the antibody produced in the animal body, and purifying the isolated antibody. The antibody may include a polyclonal and a monoclonal antibody and methods of preparing these antibodies are also known to the skilled artisan.

The following examples are provided to further illustrate the present invention but are in no way to be taken as limiting the invention. Various alterations and modifications can be made by the skilled artisan and are included within the scope of the invention.

Examples

Materials and Methods of Experiment

Rabbit chondrocyte culture

Chondrocytes were isolated from rabbit costal cartilage using, with necessary modifications, the method of Kato et al. (Kato et al.: J. Cell Biol., vol. 100, pp. 477-485, 1985). Specifically, the resting cartilage of ribs in 4-week old male Japanese albino rabbits (Hiroshima

Creating MTf overexpressing ATDC5 variant cells

Rabbit MTf cDNA (Kawamoto T. et al., EJB, 1988) of either full length or a truncated form which had the 28 residues from C-terminal necessary for GPI anchor binding
5 deleted was inserted into pcDNA3.1/Zeo(+) plasmid expression vector (containing a cytomegalovirus very early promoter/enhancer; Invitrogen, San Diego, CA). Specifically, an EcoRI-NotI fragment including the full length was excised from a vector and inserted at the EcoRI-
10 NotI site of pcDNA3.1/Zeo(+). To create a variant which lacks the GPI anchor binding site, a fragment was prepared having a stop codon inserted 28 amino acids upstream of the C-terminal and after confirming its sequence, the fragment was inserted at the EcoRI-NotI site of pcDNA3.1/Zeo(+).

15 In these ways, there were prepared a plasmid having a full length MTf cDNA (MTf Full) as an insert and a plasmid having GPI anchor-lacking MTf cDNA (MTf(-)GPI) as an insert; the two plasmids (pMTf Full and pMTf(-)GPI) were each transfected to ATDC5 cells (Riken, Tsukuba, Japan)
20 using SuperFect Transfection Reagent (QIAGEN). By selection with Zeocin (Invitrogen), stable transformants were prepared.

Specifically, 2×10^5 ATDC5 cells were seeded in 10-cm culture dishes. On the next day, 2 μ g each of the
25 plasmid DNAs to be introduced (pMTf Full and pMTf(-)GPI) and about 40 μ L of SuperFect Transfection Reagent in solution were individually dissolved in a serum-free medium and stored until use, when they were rapidly mixed together

and added to ATDC5 cells washed with a serum-free medium. After incubation at 37°C for 1 hour in the atmosphere of 5% CO₂ gas, a serum-supplemented medium was added and cultivation was conducted for an additional day. A control
5 group was prepared by transfecting only the vector.

One day after the transfection, selection was started in a serum-supplemented medium containing 50 µg/mL of Zeocin and cell culture was continued for 2 weeks with medium change effected on every third day. As a result,
10 there were obtained ATDC5 variant cell lines that would assure stable expression of MTf Full and MTf(-)GPI and these variant cell lines were subcultured in a serum-supplemented medium containing 50 µg/mL of Zeocin.

A scheme of the procedure of creating MTf
15 overexpressing ATDC5 variant cells is shown in Fig. 1.

Expression of rabbit MTf gene in ATDC5 variant cell lines

Expression of rabbit MTf gene in ATDC5 variant cell lines was confirmed by Northern blotting. Specifically, total RNA was prepared from the ATDC5 variant cell lines by
20 the guanidine thiocyanate method; 10 µg of the total RNA was electrophoresed on a 1% agarose gel containing 2.2 mol/L of formaldehyde and transferred onto Hybond-N membrane (Amersham). The membrane was hybridized with a ³²P labeled 2.2 kb rabbit MTf cDNA probe at 42°C for 16 hours.
25 After washing the membrane, a BioMax X-ray film (Kodak) was exposed to the membrane at -80°C to detect signals. The result is shown in Fig. 2.

In the MTf Full cell line, it was found that the

rabbit MTf gene have been expressed strongly in clone Nos. 1, 4 and 5.

In the MTf(-)GPI cell line, it was found that the rabbit MTf gene have been expressed strongly in clone Nos. 3, 3N, 8, 9 and 10. In the Examples, (-)GPI-3 was used.

Expression of rabbit MTf protein in ATDC5 variant cell lines

Expression of rabbit MTf protein in ATDC5 variant cell lines was confirmed by Western blotting. Specifically, membrane fraction protein was prepared from the ATDC5 variant cell lines, subjected to SDS-PAGE at 10 µg/lane, and transferred to a polyvinylidene difluoride membrane (Milipore). After the transfer, the membrane was blocked with 4% skimmed milk and reacted with anti-MTf serum [1:500 dilution; Eur. J. Biochem, 256, 503-509 (1988)] at 4°C for 14 hours, then reacted with ¹²⁵I sheep anti-mouse IgG(Fab')₂ fragment (Amersham) at room temperature for 2 hours. The membrane was washed and a BioMax X-ray film was exposed to the membrane at -80°C for analysis. The result is shown in Fig. 3.

In the MTf Full cell line, it was found that the rabbit MTf protein have been expressed strongly in clone Nos. 1 and 5. These clones were named MTf overexpressing cell lines (Full-1 and Full-5).

Example 1: Chondrogenic differentiation in MTf overexpressing cell lines

The MTf overexpressed cell lines (4.0×10^4 cells) were seeded in 6-multiwell plates and cultured in a

maintenance medium at 37°C in a 5% CO₂ gas phase.

The MTf overexpressing cell lines to be investigated were Full-1 and Full-5 which were found to have expressed both the MTf gene and protein. The MTf(-)GPI cell line to
5 be investigated was GPI-3 which was found to have expressed the MTf gene.

As control cells, ATDC5 cells and pC-1 (vector alone) were prepared in the same manner as described above and their morphological features were examined under a
10 microscope. Cell morphology was examined with an Olympus phase-contrast microscope. Two microscopic fields were taken for each culture system and at least 200 cells were counted to calculate the proportion of round cells.

In the absence of insulin, the control cells (pC-1)
15 did not differentiate to chondrocytes; on the other hand, the MTf overexpressing cell lines (Full-1 and Full-5) and the MTf(-)GPI cell line [(-)GPI-3] started to differentiate within 20 days and 29 days after seeding, almost all regions of the cells had differentiated to chondrocytes
20 (Fig. 4).

The similar test was conducted in the presence of insulin (10 µg/mL) (insulin was added at day 0). The result was the same as in the absence of insulin (Fig. 5), except that, in the presence of insulin, further
25 differentiation was induced in the MTf overexpressing cell lines, namely, more cells "rounded" like chondrocytes than in the absence of insulin.

The above results show the effectiveness of MTf in

inducing chondrogenic differentiation, which was exhibited even in the absence of insulin.

Example 2: Effect of Adding the Conditioned Medium of Rabbit Chondrocyte Culture

5 Resting rabbit chondrocytes (1×10^6 cells) were seeded in 10-cm culture dishes and cultured in medium A (10 mL) at 37°C in the atmosphere of 5% CO₂ gas. Two days after confluence, medium A was replaced by serum-free medium B (5 mL). After 24 hours of cell culture, the conditioned
10 medium (CM) was recovered and subjected to an experiment. After the recovery, CM was supplemented with fetal calf serum at a concentration of 5%.

 MTf overexpressing cell line (Full-5) and the control cell line (pC-1) were seeded at 8.0×10^4 cells in 6-
15 multiwell plates and cultured in a maintenance medium at 37°C in the atmosphere of 5% CO₂ gas. Three days after confluence (day 7), the previously recovered CM was added to give a concentration of 60% in the overall liquid culture; at the same time, 10 µg/mL of bovine insulin was
20 added. Cultivation was continued for additional 48 hours at 37°C in a 5% CO₂ gas phase.

 Forty-eight hours after the addition of CM, almost all cells of the MTf overexpressing cell line to which CM was added [Full-5(+)-CM] differentiated to chondrocytes
25 which synthesized active substrate and which resembled paving stones. The cells of the MTf expressing cell line to which no CM was added [Full-5(-)-CM] had almost the same morphology as the control cell lines in which only a vector

was expressed [pC-2(+)CM and (-)CM]. The cell lines in which only a vector was expressed had no visible induced differentiation to chondrocytes due to the addition of CM (Fig. 6).

5 These results show the presence of an MTf activating agent in CM.

Example 3: Chondrogenic differentiation due to
Overexpression of Antisense MTf RNA

ATDC5 variant cell lines (A-01, A-05, A-08, A-09, A-
10 11, A-12, A-23 and A-24) in which mouse antisense MTf RNA was overexpressed were prepared by the similar method used in preparing the ATDC5 variant cell lines in which MTf was forcibly expressed. The mouse antisense MTf RNA was prepared as follows: cDNA fragments of mouse MTf were
15 amplified by PCR (using primers 5'-GGTGTGTTGAGGGGCGTGGACTCT-3' (SEQ ID NO: 9) and 5'-TCACCAACGGCTTTGAGCACATCAC-3' (SEQ ID NO: 10), inserted into pGEM-T Easy Vector (Promega), excised with ApaI-NotI and inserted into pCDNA3.1/Zeo(+) at ApaI-NotI site (i.e.,
20 inserted in reverse direction). The sequence of the inserted portion is identified by SEQ ID NO: 11.

A control group was also prepared by transfecting only the vector (pC-1).

The expression of mouse MTf antisense was examined by
25 Northern blotting using the above-mentioned ApaI-NotI fragment as a probe.

A criterion for the suppression of chondrogenic differentiation was the suppression of synthesis of a

cartilage proteoglycan, aggrecan, and a test was conducted by RT-PCR Southern blotting both in the presence of insulin (added in an amount of 10 µg/mL after day 4) and in its absence. Specifically, total RNA was extracted from the
5 cells of each clone by the guanidine thiocyanate method; single-stranded cDNA was synthesized from the extracted total RNA (1 µg) using SUPERScript pre-amplification system kit (Life Technologies); using the cDNA as a template, PCR was performed on the aggrecan gene with a pair of primers
10 5'-TGCTACTTCATCGACCC-3' (forward) (SEQ ID NO: 12) and 5'-AAAGACCTCCCCTCCATCT-3' (reverse) (SEQ ID NO: 13); the PCR reaction mixture was electrophoresed on a 1% agarose gel and transferred onto Hybond-N membrane (Amersham). The membrane was hybridized with a ³²P labeled antisense MTF
15 probe and mouse aggrecan cDNA probe at 42°C for 16 hours. The membrane was washed with 0.2 x SSC which contains 0.5% SDA and a BioMax X-ray film (Kodak) was exposed to the membrane at -80°C to detect signals.

The result is shown in Fig. 7. Antisense MTF RNA was
20 expressed most strongly in variant cell line A-12 (lane 6), followed by A-11 (lane 5) in strength. Correspondingly, expression of the aggrecan gene was suppressed most effectively in variant cell line A-12 (lane 6), followed by A-11 (lane 5) in effectiveness. While expression of the
25 aggrecan gene was suppressed in the absence of insulin, it was more effectively suppressed in the presence of insulin.

CLAIMS

1. A cartridge formation stimulator containing a membrane-bound transferrin-like protein (MTf).
2. The chondrogenesis stimulator according to claim 1, wherein MTf is selected from the group consisting of rabbit p76 protein, human p97 protein, and a protein demonstrating the MTf activity that has an amino acid sequence encoded by DNA which hybridizes, under stringent conditions, with a DNA coding for p76 protein or p97 protein.
3. The chondrogenesis stimulator according to claim 1, wherein the MTf is selected from the following:
 - 1) a protein having the amino acid sequence of SEQ ID NO: 2;
 - 2) a protein having the amino acid sequence of SEQ ID NO: 4;
 - 3) a protein having the amino acid sequence of SEQ ID NO: 15; and
 - 4) a protein demonstrating the MTf activity that has an amino acid sequence encoded by DNA which hybridizes, under stringent conditions, with a DNA encoding the protein of SEQ ID NO: 2, 4 or 15.
4. The chondrogenesis stimulator according to claim 2, wherein the MTf is human p97 protein.
5. A chondrogenesis stimulator containing soluble MTf.
6. The chondrogenesis stimulator according to claim 5, wherein the soluble MTf lacks the GPI anchor region.
7. An agent for gene therapy to promote chondrogenesis which contains as an active ingredient an expression vector

incorporating a DNA encoding any one of the following proteins:

- 1) a protein having the amino acid sequence of SEQ ID NO: 2;
- 2) a protein having the amino acid sequence of SEQ ID NO: 4;
- 3) a protein having the amino acid sequence of SEQ ID NO: 15;
- 4) a protein demonstrating the MTF activity that has an amino acid sequence encoded by DNA which hybridizes, under stringent conditions, with a DNA coding for the protein of SEQ ID NO: 2, 4 or 15; and

5) a protein which is the same as protein 1), 2), 3) or 4), except that it lacks the GPI anchor region.

8. The chondrogenesis stimulator according to claim 1 which is used in combination with an MTF activating agent.

9. The chondrogenesis stimulator according to claim 1 which is used in combination with insulin or an insulin-like growth factor.

10. The chondrogenesis stimulator according to any one of claims 1 - 9 for treating at least one bone disease selected from the following diseases in which chondrogenic differentiation is involved: OA (osteoarthritis); RA (rheumatoid arthritis); injury of articular cartilage due to trauma; maintenance of chondrocyte phenotypes in autologous chondrocyte transplantation; reconstruction of cartilage in the ear, trachea or nose; osteochondritis dissecans; regeneration of intervertebral disk or meniscus; bone fracture; and osteogenesis from cartilage.

11. A chondrogenic differentiation suppressing agent containing an MTf antagonist.

12. The chondrogenic differentiation suppressing agent according to claim 11, wherein the MTf antagonist is an anti-MTf antibody or an oligonucleotide or an oligonucleotide analog that are hybridizable with a nucleic acid encoding MTf.

13. A method for screening an MTf activating agent which comprises the steps of:

1) preparing a cell line in which MTf is overexpressed, wherein said cell line retains the ability to differentiate to chondrocytes but hardly differentiate without stimulation;

2) adding candidate substances to the cell line prepared in step 1) and culturing it for a specified period of time; and

3) examining the cell line for induced chondrogenic differentiation and selecting an MTf activating agent from the candidate substances.

14. An MTf activating agent obtained by the method according to claim 13.

15. A chondrogenesis stimulator containing an MTf activating agent obtained by the method according to claim 13.


16. MTf which lacks the GPI anchor region.

ABSTRACT

There are provided a chondrogenesis stimulator containing MTf, a chondrogenic differentiation suppressing agent containing an MTf antagonist, a screening method for
5 obtaining an MTf activating agent, an MTf activating agent obtained by the screening method, a chondrogenesis stimulator containing an MTf activating agent as obtained by the screening method, and MTf which lacks the GPI anchor region.

Fig. 1

Construction of expression vector (pc-DNA 3.1 (+) plasmid)



Constructed expression vectors

MTf Full



MTf (-) GPI



Stable transfection



Checking for the expression of MTf mRNA by Northern blotting

Checking for the expression of MTf protein by Western blotting

Fig.2




Fig.3





Fig.4

Fig.5







Fig.6

*Fig.7***Antisense**

1 2 3 4 5 6 7 8

**MTf mRNA****day 23 (-) insulin****AggreCan****day 17 (+) insulin****AggreCan**

1 2 3 4 5 6 7 8 9

- 1 = A-01
- 2 = A-05
- 3 = A-08
- 4 = A-09
- 5 = A-11
- 6 = A-12
- 7 = A-23
- 8 = A-24
- 9 = pC-1

100499571021902
 10/049957
 Rec'd PCT/PTO 05 JUL 2002

SEQUENCE LISTING

<110> KATO, Yukio
 FUJIMOTO, Katsumi

<120> CHONDROGENESIS PROMOTERS

<130> KATO=21

<140> 10/049,957

<141> 2002-02-19

<150> PCT/JS00/05590

<151> 2000-08-21

<150> JP 232966/1999

<151> 1999-08-19

<160> 15

<170> PatentIn version 3.1

<210> 1

<211> 2388

<212> DNA

<213> Oryctolagus cuniculus

<400> 1

gccgccgctc actcggttcgc actcggactc agaccagtc cgacccctg gactgcgcca	60
tgcggtgccg aagcgcggct atgtggatct tcctggccct gcgcaccgca ctcggcagcg	120
tggagggtgcg gtgggtgcacc gcgtccgagc ccgagcagca gaagtgcgag gacatgagcc	180
aggccttccg cgaagccggc ctccagcccc cctgctgtg cgtgcagggc acctcgcccg	240
accactgcgt ccagctcatc gcggcccacg aggccgacgc catcactctg gacggaggag	300
ccatttacga ggcggggaag gaacacggcc tgaagcccg ggtgggagaa gtgtatgacc	360
aagagggtggg cacctcctac tacgtgtgtg ccgtgggtcaa gaggagctcc aacgtgacca	420
tcaacaccct gagaggcgtg aagtcctgcc acacgggcat caaccgcacg gtgggctgga	480
acgtgcctgt gggctacctg gtggacagcg gccgcctctc agtgatgggc tgtgacgtgc	540
tcaaagcggc cagcgagtac ttccggggga gctgcgtccc tggggcagga gagaccagat	600
actcggagtc cctctgtcgc ctctgccggg gcgacacctc cggggagggg gtgtgtgaca	660
agagccccct ggagcgggtac tacgactaca gcggggcctt ccggtgcctg gcagaaggcg	720
caggggacgt ggcctttgtg aagcacagca ccgtgctgga gaacacggat gggagaacac	780
tgcctcctg gggccacatg ctgatgtcac gggactttga gctgctgtgc cgggacggca	840
gccggggcag cgtcaccgag tggcagcact gccacctggc ccgggtgccc gccacgccc	900

tgggtggtcctg	ggccgacacc	gacgcaggcc	tcattcttccg	gcttctcaat	gagggccagc	960
ggctgttcag	ccacgagggc	agcagcttcc	agatgttcag	ctcggaggcc	tacggccaga	1020
agaacctgct	gttcaaagac	tccacgctgg	agctgggtgcc	catcgccaca	cagacctacg	1080
aggcctggct	gggccccgag	tacctgcacg	ccatgaaggg	tctgctctgt	gaccccaacc	1140
ggctgcccc	atacctgcgc	tgggtgcgtgc	tgtccacccc	cgagatccag	aagtgtggag	1200
acatggccgt	ggccttcagc	cggcagaggc	tcaagccgga	gatccagtgt	gtctcggcgg	1260
agtcccccca	gcactgcatg	gagcagatcc	aggctgggca	catcgatgct	gtgacctga	1320
acggggagga	cattcacaca	gcggggaaga	cttatgggct	gatcccggt	gcgcggggagc	1380
tgtatgccgc	ggacgacagg	agtaactcgt	acttcgtggg	ggccgtgggtg	aagcgagaca	1440
gcgcctacgc	cttcaccgtg	gacgagctgc	gcgggaagcg	ctcctgccac	cccggcttcg	1500
gcagcccggc	cggctgggac	gtcccgggtg	gcgcctcat	ccactggggc	tacatccggc	1560
ccaggaactg	cgacgtcctc	acagcgggtg	gtcagttctt	caacgccagc	tgtgtgccgg	1620
tgaacaaccc	caagaagtac	ccctcctcgc	tgtgcgcact	ctgcgtgggt	gacgagcagg	1680
gccgcaacaa	gtgcactggc	aacagccagg	agcgggtacta	tggcgacagt	ggcgccttca	1740
gggtgcctggt	ggaggggtgca	ggggacgtgg	ccttcgtcaa	gcacacgacc	atctttgaca	1800
acacaaatgg	ccacaatccc	gagccgtggg	ctgcccattct	gaggagccag	gactacgagc	1860
tgctgtgccc	caacggcgcg	cgagctgagg	cgcaccagtt	tgccgcctgc	aacctggccc	1920
agattccgtc	ccacgcgcgc	atgggtgcggc	ccgacaccaa	catcttcacc	gtttacggac	1980
tgctggacaa	ggcccaggac	ctgttttgag	acgaccacaa	caagaacggg	ttcaagatgt	2040
tcgactcctc	cagctaccac	ggccgagacc	tgtctttcaa	ggacgccacg	gtgcgcgctg	2100
tgccctgtggg	cgagaggacc	acctaccagg	actggctggg	gccggactac	gtggcggctc	2160
tggaagggat	gcagtcacag	cgggtgctcag	gggcagccgt	cggcgcccc	ggggcctcgc	2220
tgctgcgcgt	gctgccccctg	gctgcggggc	tctgtctgtc	ttcgtcttga	gagcagcccc	2280
gggcagcctc	ggccccggca	ggggagcctg	cgcggaagct	tctgaacga	gcccgcgccc	2340
tggtgtgatg	tggttacctc	ggcgagccgc	ggggccgcgc	ttcccccg		2388

Met	Arg	Cys	Arg	Ser	Ala	Ala	Met	Trp	Ile	Phe	Leu	Ala	Leu	Arg	Thr	1	5	10	15
Ala	Leu	Gly	Ser	Val	Glu	Val	Arg	Trp	Cys	Thr	Ala	Ser	Glu	Pro	Glu	20	25	30	
Gln	Gln	Lys	Cys	Glu	Asp	Met	Ser	Gln	Ala	Phe	Arg	Glu	Ala	Gly	Leu	35	40	45	
Gln	Pro	Ala	Leu	Leu	Cys	Val	Gln	Gly	Thr	Ser	Ala	Asp	His	Cys	Val	50	55	60	
Gln	Leu	Ile	Ala	Ala	His	Glu	Ala	Asp	Ala	Ile	Thr	Leu	Asp	Gly	Gly	65	70	75	80
Ala	Ile	Tyr	Glu	Ala	Gly	Lys	Glu	His	Gly	Leu	Lys	Pro	Val	Val	Gly	85	90	95	
Glu	Val	Tyr	Asp	Gln	Glu	Val	Gly	Thr	Ser	Tyr	Tyr	Ala	Val	Ala	Val	100	105	110	
Val	Lys	Arg	Ser	Ser	Asn	Val	Thr	Ile	Asn	Thr	Leu	Arg	Gly	Val	Lys	115	120	125	
Ser	Cys	His	Thr	Gly	Ile	Asn	Arg	Thr	Val	Gly	Trp	Asn	Val	Pro	Val	130	135	140	
Gly	Tyr	Leu	Val	Asp	Ser	Gly	Arg	Leu	Ser	Val	Met	Gly	Cys	Asp	Val	145	150	155	160
Leu	Lys	Ala	Val	Ser	Glu	Tyr	Phe	Gly	Gly	Ser	Cys	Val	Pro	Gly	Ala	165	170	175	
Gly	Glu	Thr	Arg	Tyr	Ser	Glu	Ser	Leu	Cys	Arg	Leu	Cys	Arg	Gly	Asp	180	185	190	
Thr	Ser	Gly	Glu	Gly	Val	Cys	Asp	Lys	Ser	Pro	Leu	Glu	Arg	Tyr	Tyr	195	200	205	
Asp	Tyr	Ser	Gly	Ala	Phe	Arg	Cys	Leu	Ala	Glu	Gly	Ala	Gly	Asp	Val	210	215	220	
Ala	Phe	Val	Lys	His	Ser	Thr	Val	Leu	Glu	Asn	Thr	Asp	Gly	Arg	Thr	225	230	235	240
Leu	Pro	Ser	Trp	Gly	His	Met	Leu	Met	Ser	Arg	Asp	Phe	Glu	Leu	Leu	245	250	255	
Cys	Arg	Asp	Gly	Ser	Arg	Ala	Ser	Val	Thr	Glu	Trp	Gln	His	Cys	His	260	265	270	
Leu	Ala	Arg	Val	Pro	Ala	His	Ala	Val	Val	Val	Arg	Ala	Asp	Thr	Asp	275	280	285	
Ala	Gly	Leu	Ile	Phe	Arg	Leu	Leu	Asn	Glu	Gly	Gln	Arg	Leu	Phe	Ser	290	295	300	
His	Glu	Gly	Ser	Ser	Phe	Gln	Met	Phe	Ser	Ser	Glu	Ala	Tyr	Gly	Gln				

[illegible]

His Ala Val Met Val Arg Pro Asp Thr Asn Ile Phe Thr Val Tyr Gly
625 630 635 640

Leu Leu Asp Lys Ala Gln Asp Leu Phe Gly Asp Asp His Asn Lys Asn
645 650 655

Gly Phe Lys Met Phe Asp Ser Ser Ser Tyr His Gly Arg Asp Leu Leu
660 665 670

Phe Lys Asp Ala Thr Val Arg Ala Val Pro Val Gly Glu Arg Thr Thr
675 680 685

Tyr Gln Asp Trp Leu Gly Pro Asp Tyr Val Ala Ala Leu Glu Gly Met
690 695 700

Gln Ser Gln Arg Cys Ser Gly Ala Ala Val Gly Ala Pro Gly Ala Ser
705 710 715 720

Leu Leu,Pro Leu Leu Pro Leu Ala Ala Gly Leu Leu Leu Ser Ser Leu
725 730 735

```
<210> 3
<211> 2368
<212> DNA
<213> Homo sapiens
```

<400>	3						
gcggacttcc	tgggacccgg	accagacccc	agcccgcccc	cagccagccc	cgacggcgcc		60
atgcgggggtc	cgagcgggggc	tctgtggctg	ctcctggctc	tgcgcaccgt	gctcggaggc		120
atggaggtgc	ggtggtgctc	cacctcggac	ccagagcagc	acaagtgcgg	caacatgagc		180
gaggccttcc	gggaagcggg	catccagccc	tccctcctct	gcgtccgggg	cacctccgcc		240
gaccactgct	tccagctcat	cgcggcccag	gaggctgacg	ccatcactct	ggatggagga		300
gccatctatg	aggcgggaaa	ggagcacggc	ctgaagccgg	tggtagggcg	agtgtacgat		360
caagaggctc	gtacctccta	ttacgcctgt	gctgtggtca	ggaggagctc	ccatgtgacc		420
attgacaccc	tgaaaggcgt	gaagtccctg	cacacggggc	tcaatcgcac	agtgggctgg		480
aacgtgccc	tgggctacct	ggtggagagc	ggccgcctct	cggatgatgg	ctgcgatgta		540
ctcaaagctg	tcagcgacta	ttttgggggc	agctgcgtcc	cgggggcagg	agagaccagt		600
tactctgagt	ccctctgtcg	cctctgcagg	ggtgacagct	ctggggaagg	ggtgtgtgac		660
aagagcccc	tggagagata	ctacgactac	agcggggcct	tccggtgcct	ggcggaaggg		720
gcaggggacg	tggtttttgt	gaagcacagc	acggtactgg	agaacacgga	tgggaagacg		780
cttccctcct	ggggccaggc	cctgctgtca	caggacttcg	agctgctgtg	ccgggatggt		840
agccggggcc	atgtcaccca	gtggaggcag	tgccatctgg	cccgggtgcc	tgtcacgcc		900

gtggtggtcc	gggccgacac	agatgggggc	ctcatcttcc	ggctgctcaa	cgaaggccag	960
cgtctgttca	gccacgaggg	cagcagcttc	cagatgttca	gctctgaggg	ctatggccag	1020
aaggatctac	tcttcaaaga	ctctacctcg	gagcttgtgc	ccatcgccac	acagacctat	1080
gaggcgtggc	tggggccatga	gtacctgcac	gccatgaagg	gtctgctctg	tgaccccaac	1140
cggctgcccc	cctacctgcg	ctggtgtgtg	ctctccactc	ccgagatcca	gaagtgtgga	1200
gacatggccg	tggccttccg	ccggcagcgc	ctcaagccag	agatccagtg	cgtgtcagcc	1260
aagtcccccc	aacactgcat	ggagcggatc	caggctgagc	aggtegacgc	tgtgacccta	1320
agtggcgagg	acattttacac	ggcggggaag	aagtaaggcc	tggttcccgc	agccggcgag	1380
cactatgccc	cggaagacag	cagcaactcg	tactacgtgg	tggccgtggg	gagacggggac	1440
agctcccacg	ccttcacctt	ggatgagctt	cggggcaagc	gctcctgcca	cgccggtttc	1500
ggcagccctg	caggctggga	tgtccccgtg	ggtgccctta	ttcagagagg	cttcatccgg	1560
cccaaggact	gtgacgtect	cacagcagtg	agcgagttct	tcaatgccag	ctgcgtgccc	1620
gtgaacaacc	ccaagaacta	ccccctctcg	ctgtgtgcac	tgtgcgtggg	ggacgagcag	1680
ggccgcaaca	agtgtgtggg	caacagccag	gagcggtatt	acggctaccg	cggcgccctc	1740
aggtgcctgg	tggagaatgc	gggtgacgtt	gccttcgtca	ggcacacaac	cgtctttgac	1800
aacacaaaac	gccacaatte	cgagccctgg	gctgctgagc	tcaggtcaga	ggactatgaa	1860
ctgctgtgcc	ccaacggggc	ccgagccgag	gtgtcccagt	ttgcagcctg	caacctggca	1920
cagataccac	cccacgccgt	gatggtcggg	cccgcacca	acatcttcac	cgtgtatgga	1980
ctgctggaca	aggcccagga	cctgttttga	gacgaccaca	ataagaacgg	gttcaaaatg	2040
ttcgactcct	ccaactatca	tggccaagac	ctgcttttca	aggatgccac	cgtccggggc	2100
gtgcctgtcg	gagagaaaaac	cacctaccgc	ggctggctgg	ggctggacta	cgtggcgggc	2160
ctggaagggg	tgtcgtctca	gcagtgctcg	ggcgcagcgg	ccccggcgcc	cggggcgccc	2220
ctgctccccg	tgtctgtgcc	cgcctctgcc	gcccgcctgc	tcccgcctcg	cctctgagcc	2280
cggccgcccc	gccccagagc	tccgatgcc	gcccggggag	ttccgcgggc	ggcctctcgc	2340
gctgcggaat	ccagaaggaa	gctcgcga				2368

```
<210> 4
<211> 738
<212> PRT
<213> Homo sapiens
```

<400> 4

Met Arg Gly Pro Ser Gly Ala Leu Trp Leu Leu Leu Ala Leu Arg Thr
 1 5 10 15
 Val Leu Gly Gly Met Glu Val Arg Trp Cys Ala Thr Ser Asp Pro Glu
 20 25 30
 Gln His Lys Cys Gly Asn Met Ser Glu Ala Phe Arg Glu Ala Gly Ile
 35 40 45
 Gln Pro Ser Leu Leu Cys Val Arg Gly Thr Ser Ala Asp His Cys Val
 50 55 60
 Gln Leu Ile Ala Ala Gln Glu Ala Asp Ala Ile Thr Leu Asp Gly Gly
 65 70 75 80
 Ala Ile Tyr Glu Ala Gly Lys Glu His Gly Leu Lys Pro Val Val Gly
 85 90 95
 Glu Val Tyr Asp Gln Glu Val Gly Thr Ser Tyr Tyr Ala Val Ala Val
 100 105 110
 Val Arg Arg Ser Ser His Val Thr Ile Asp Thr Leu Lys Gly Val Lys
 115 120 125
 Ser Cys His Thr Gly Ile Asn Arg Thr Val Gly Trp Asn Val Pro Val
 130 135 140
 Gly Tyr Leu Val Glu Ser Gly Arg Leu Ser Val Met Gly Cys Asp Val
 145 150 155 160
 Leu Lys Ala Val Ser Asp Tyr Phe Gly Gly Ser Cys Val Pro Gly Ala
 165 170 175
 Gly Glu Thr Ser Tyr Ser Glu Ser Leu Cys Arg Leu Cys Arg Gly Asp
 180 185 190
 Ser Ser Gly Glu Gly Val Cys Asp Lys Ser Pro Leu Glu Arg Tyr Tyr
 195 200 205
 Asp Tyr Ser Gly Ala Phe Arg Cys Leu Ala Glu Gly Ala Gly Asp Val
 210 215 220
 Ala Phe Val Lys His Ser Thr Val Leu Glu Asn Thr Asp Gly Lys Thr
 225 230 235 240
 Leu Pro Ser Trp Gly Gln Ala Leu Leu Ser Gln Asp Phe Glu Leu Leu
 245 250 255
 Cys Arg Asp Gly Ser Arg Ala Asp Val Thr Glu Trp Arg Gln Cys His
 260 265 270
 Leu Ala Arg Val Pro Ala His Ala Val Val Val Arg Ala Asp Thr Asp
 275 280 285
 Gly Gly Leu Ile Phe Arg Leu Leu Asn Glu Gly Gln Arg Leu Phe Ser
 290 295 300

His Glu Gly Ser Ser Phe Gln Met Phe Ser Ser Glu Ala Tyr Gly Gln
305 310 315 320

Lys Asp Leu Leu Phe Lys Asp Ser Thr Ser Glu Leu Val Pro Ile Ala
325 330 335

Thr Gln Thr Tyr Glu Ala Trp Leu Gly His Glu Tyr Leu His Ala Met
340 345 350

Lys Gly Leu Leu Cys Asp Pro Asn Arg Leu Pro Pro Tyr Leu Arg Trp
355 360 365

Cys Val Leu Ser Thr Pro Glu Ile Gln Lys Cys Gly Asp Met Ala Val
370 375 380

Ala Phe Arg Arg Gln Arg Leu Lys Pro Glu Ile Gln Cys Val Ser Ala
385 390 395 400

Lys Ser Pro Gln His Cys Met Glu Arg Ile Gln Ala Glu Gln Val Asp
405 410 415

Ala Val Thr Leu Ser Gly Glu Asp Ile Tyr Thr Ala Gly Lys Lys Tyr
420 425 430

Gly Leu Val Pro Ala Ala Gly Glu His Tyr Ala Pro Glu Asp Ser Ser
435 440 445

Asn Ser Tyr Tyr Val Val Ala Val Val Arg Arg Asp Ser Ser His Ala
450 455 460

Phe Thr Leu Asp Glu Leu Arg Gly Lys Arg Ser Cys His Ala Gly Phe
465 470 475 480

Gly Ser Pro Ala Gly Trp Asp Val Pro Val Gly Ala Leu Ile Gln Arg
485 490 495

Gly Phe Ile Arg Pro Lys Asp Cys Asp Val Leu Thr Ala Val Ser Glu
500 505 510

Phe Phe Asn Ala Ser Cys Val Pro Val Asn Asn Pro Lys Asn Tyr Pro
515 520 525

Ser Ser Leu Cys Ala Leu Cys Val Gly Asp Glu Gln Gly Arg Asn Lys
530 535 540

Cys Val Gly Asn Ser Gln Glu Arg Tyr Tyr Gly Tyr Arg Gly Ala Phe
545 550 555 560

Arg Cys Leu Val Glu Asn Ala Gly Asp Val Ala Phe Val Arg His Thr
565 570 575

Thr Val Phe Asp Asn Thr Asn Gly His Asn Ser Glu Pro Trp Ala Ala
580 585 590

Glu Leu Arg Ser Glu Asp Tyr Glu Leu Leu Cys Pro Asn Gly Ala Arg
595 600 605

Ala Glu Val Ser Gln Phe Ala Ala Cys Asn Leu Ala Gln Ile Pro Pro

<400> 7
agagggactc cgagtatctg gtctc 25

<210> 8
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic

<400> 8
gtccggcccg acaccaacat cttc 24

<210> 9
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic

<400> 9
ggtgtgttga gggcggtgga ctct 24

<210> 10
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic

<400> 10
tcaccaacgg ctttgagcac atcac 25

<210> 11
<211> 614
<212> DNA
<213> Mus sp.

<400> 11
tcaccaacgg ctttgagcac atcacagccc atcactgaca gatggccgct ctctacgagg 60
taaccgacag gcacgttcca gccacagtc cggttaatgc ctgtgtggca ggacttgacg 120
cccttcaggg tgttgatggt aacattggaa ttctctctga ccacagccac ggcataatag 180
gaagtcccaa tgtcttggtc atagacttcc cccaccactg gcttcaggcc gtgtctcttc 240
cctgcctcat agatggcccc tccatccagg gtgatggcat ctgctttttg ttccttgatg 300
agctggacac agtggtcagc ggagttgccc tggacgcaga gaagggaagg acgaatgcc 360

gctccctgga aggcctcgct catgtctttg cacttctgct gctctgcgtc tgagatggta 420
 caccactgca cctccatcac acagacgaca gtgcgcaggg acaggagtag ccaaaaagtc 480
 acgctcagga gcctcatggc aacgttgggt tggctgggggt gctggcgggt ctgtcctggc 540
 ttctcttccc ctggtctctc tggccttcac tattaagcg cagcccgggg agagtccacg 600
 cccctcaaca cacc 614

<210> 12
 <211> 17
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic

<400> 12 17
 tgctacttca tcgaccc

<210> 13
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic

<400> 13 19
 aaagacctcc cctccatct

<210> 14
 <211> 4158
 <212> DNA
 <213> Mus sp.

<400> 14 60
 ggtgtgttga ggggcgtgga ctctccccgg gctgcgetta aatagtgaag gccagagaga
 ccagggaaga ggaagccagg acagaccgc cagcacccca gccaacccaa cgttgccatg 120
 aggctcctga gcgtgacttt ttggctactc ctgtccctgc gcactgtcgt ctgtgtgatg 180
 gaggtgcagt ggtgtaccat ctacagcgca gagcagcaga agtgcaaaga catgagcgag 240
 gccttccagg gagctggcat tcgtccttcc ctctctctgcg tccagggcaa ctccgctgac 300
 cactgtgtcc agctcatcaa ggaacaaaaa gcagatgcc aaccctgga tggagggggc 360
 atctatgagg cagggaagga gcaaggcctg aagccagtgg tgggggaagt ctatgaccaa 420
 gacattggga ctctctatta tgccgtggct gtggtcagga ggaattccaa tgttaccatc 480

aacaccctga agggcgctcaa gtccctgccac acaggcatta accggactgt gggctggaac 540
 gtgcctgtcg gttacctcgt agagagcggc catctgtcag tgatgggctg tgatgtgctc 600
 aaagccgttg gtgattatatt tggaggcagc tgtgtccctg gaacaggaga aaccagccat 660
 tccgagtcct tctgtcgctt ctgccgtggc gactcttctg ggcacaatgt gtgtgacaag 720
 agtcccctag agagatacta cgactacagt ggagccttcc ggtgcctggc ggaaggagcc 780
 ggtgacgtgg ccttcgtgaa gcacagcaca gtgctggaaa atactgatgg aaacaccctg 840
 ccttctctgg gcaagtccct gatgtcagag gacttcagc tactatgcag ggatggcagc 900
 cgagccgaca tcaactgagt gagacgttgc cacctggcca aggtgcctgc tcatgtgtg 960
 gtggtcaggg gtgacatgga tggcggtctc atattccaac tgctcaacga aggccagctt 1020
 ctgttcagcc acgaagacag cagcttcag atgttcagct ccaaagccta cagccagaag 1080
 aacttgetgt tcaaagactc caccttggag cttgtgccca ttgccacaca gaactacgag 1140
 gcctggctgg gccaggaata cctgcaggcc atgaaggggc tcctctgtga tcccaaccgg 1200
 ctgccccact acctgcgctg gtgtgtgctg tcagcgcccg agatccagaa gtgtggagat 1260
 atggctgtgg ccttcagccg ccagaatctc aagccggaaa ttcagtgtgt gtcggccgag 1320
 tcccctgagc actgcatgga gcagatccag gctgggcaca ctgacgctgt gactctgagg 1380
 ggcgaggaca ttacagggc aggaaagggt tacggcctgg ttccggcggc cggggagctg 1440
 tatgtgagg aggacaggag caattcctac tttgtggtgg ctgtggcaag aagggacagc 1500
 tctactcct tcaccctgga cgagcttcgc ggcaagcgtt cctgccacc ctaactgggc 1560
 agcccagcgg gctgggaggt gcccatcggc tccctcatcc agcggggctt catccggccc 1620
 aaggactgtg atgtcctcac agcggtgagc cagttcttca atgccagctg cgtgcctgtc 1680
 aacaacccta agaactacc ttccgcacta tgtgcgctct gcgtgggaga cgagaagggc 1740
 cgcaacaaat gtgtggggag cagccaggag agatactacg gctacagcgg ggccttcagg 1800
 tgccttgtgg agcatgcagg ggacgtggct ttcgtcaagc acacgactgt ctttgagaac 1860
 acaaatggtc acaatcctga gccttgggt tctcacctca ggtggcaaga ctatgaacta 1920
 ctgtgcccc aatggggcag ggctgaggt gaccagttcc aagcttgcaa cctggcacia 1980
 atgccatccc acgtgtcat ggtccgtcca gacaccaaca tcttactgt gtatggactt 2040
 ctggacaagg cccaggacct gtttgagac gaccataaca agaacggttt ccaaagtgtt 2100
 gactcctcca aatatcacag ccaagacct cttttcaaag atgtacagt ccgagcgggt 2160
 ccagtcgggg agaaaaccac atacctggac tggctgggtc ctgactatgt ggttgcgctg 2220

gaggggatgt tgtctcagca gtgctccggt gcaggggccg cgggtgcagcg agtccccctg 2280
 ctggccctgc tectgctgac cctggctgca ggccctcttc ctgcgcttct ctgaagaccg 2340
 ctgcttcagg ccacgcccag agcagggaaa gctacagagc tcaaccggaa gaaaccagga 2400
 catcagctaa ccctgcagga gagcgcgggg cgggatgagg agaggcaagg tgagaactca 2460
 cacacacaca caagcctccg aggtgcgatt ctaacccaaa gagaaatttc tagaatcagg 2520
 atgattgtta aggccaagtc tccccacttg ctggagccct caatacctga ggcgactggc 2580
 gtagtagcca gtcactctc ccacaccggt ggcgccagca gcgaacctgt gcctcccacc 2640
 tggagcctcc tggttggtg ggggtggttaa gggggggggg gggagagtga agatgctggt 2700
 tgccatggca accgtggagc agcttcacgc ctctgtaccg gccacctggt gagatgccaa 2760
 ggaaggagca caccaccaac ctagggaacc tgtgcgacac actaccaccc agcagcccct 2820
 gctctcgtg cccaccgct ctctctatg ggcactgtc caccaaggcc acaccgtcgg 2880
 aggggcaagg ctgctgagca catcagcctt ctgatgtgac accaaccaag gagcccagcc 2940
 ctctggacag caagattttg ctgactggg atgggaggaa ggccagagct gtactgtggg 3000
 gatgaagtcc tccaaaacc tcagaggaag gaagtgcgcc caccttccca ttaagaatgt 3060
 tagtgtgtga gaaacttgat gcaggggtga aactatcctg tttaacggct cccgtggcaa 3120
 gcaggacttg cgctgtctgc gctgcctgga cctcactgca caatgaaact gttgccgaga 3180
 ttctattgtt tgctctcctg gtctcagtct caacattagt tttctcctg ccttcatata 3240
 ccccttccca catcaccag caagcacgca cgcgcacag cacacgcaca caccttatcc 3300
 gtgtgaacat atctgaacat atctgcttgt ctgaagaagt aggagctaac caaaaataac 3360
 ttctgtcat gagctgggcc ttgggatata ccacgagcca ggggattggg gagagccctg 3420
 tcttcccttc accctgcacc tggtgggcag ttgcatcttt cgagaggatc cctggttctc 3480
 tcgaactgtg agagccaagg cctaggctgc cattttgcca ttgttctctc gagaaccaga 3540
 aaaagttttc caaagctacc agctcttacc ccagatcttg ttcccttaaa aaaaagtaat 3600
 aaataaaaag gagaagaaac aggagcaaac agccatcgtc agcacactgg aagcagcgtg 3660
 ggccgggagc tatttgtgtc ttggtctgtg tggggggcct cagatcccaa tgacaggcca 3720
 ggttcccagt ggctgcgcc cactgtggg cgacgacggg acagatcctt tccatggctc 3780
 accagtagag aaggtcctgg cagtgtccca gccagagtca cacaatcctg aggaaaatcg 3840
 gtcaccatgg tgcttgggag agcaagcccc tctcctccc agtacacagc catccattct 3900
 tctctgagct ggggacttca cagtgagaag tgtactctgt gtgggcgact gtgctgccca 3960

aagtgtgatg tctgtgccgt gtgcctttca ggtgtgactt tgaagagcgt tgtgtaaatg 4020
 acgtctgatt gccatggggcc actgctgtgt ttgtgctaaa gaaagacatt ggtttctttt 4080
 taaaataaag ccatatatcc ctgcatacgc agaggcttgg atcctgggtgg aaaaaaaaaa 4140
 aaaaaaaaaa aaaaaaaaaa 4158

<210> 15
 <211> 738
 <212> PRT
 <213> Mus sp.

<400> 15

Met Arg Leu Leu Ser Val Thr Phe Trp Leu Leu Leu Ser Leu Arg Thr
 1 5 10 15
 Val Val Cys Val Met Glu Val Gln Trp Cys Thr Ile Ser Asp Ala Glu
 20 25 30
 Gln Gln Lys Cys Lys Asp Met Ser Glu Ala Phe Gln Gly Ala Gly Ile
 35 40 45
 Arg Pro Ser Leu Leu Cys Val Gln Gly Asn Ser Ala Asp His Cys Val
 50 55 60
 Gln Leu Ile Lys Glu Gln Lys Ala Asp Ala Ile Thr Leu Asp Gly Gly
 65 70 75 80
 Ala Ile Tyr Glu Ala Gly Lys Glu His Gly Leu Lys Pro Val Val Gly
 85 90 95
 Glu Val Tyr Asp Gln Asp Ile Gly Thr Ser Tyr Tyr Ala Val Ala Val
 100 105 110
 Val Arg Arg Asn Ser Asn Val Thr Ile Asn Thr Leu Lys Gly Val Lys
 115 120 125
 Ser Cys His Thr Gly Ile Asn Arg Thr Val Gly Trp Asn Val Pro Val
 130 135 140
 Gly Tyr Leu Val Glu Ser Gly His Leu Ser Val Met Gly Cys Asp Val
 145 150 155 160
 Leu Lys Ala Val Gly Asp Tyr Phe Gly Gly Ser Cys Val Pro Gly Thr
 165 170 175
 Gly Glu Thr Ser His Ser Glu Ser Leu Cys Arg Leu Cys Arg Gly Asp
 180 185 190
 Ser Ser Gly His Asn Val Cys Asp Lys Ser Pro Leu Glu Arg Tyr Tyr
 195 200 205
 Asp Tyr Ser Gly Ala Phe Arg Cys Leu Ala Glu Gly Ala Gly Asp Val
 210 215 220

Ala Phe Val Lys His Ser Thr Val Leu Glu Asn Thr Asp Gly Asn Thr
225 230 235 240

Leu Pro Ser Trp Gly Lys Ser Leu Met Ser Glu Asp Phe Gln Leu Leu
245 250 255

Cys Arg Asp Gly Ser Arg Ala Asp Ile Thr Glu Trp Arg Arg Cys His
260 265 270

Leu Ala Lys Val Pro Ala His Ala Val Val Val Arg Gly Asp Met Asp
275 280 285

Gly Gly Leu Ile Phe Gln Leu Leu Asn Glu Gly Gln Leu Leu Phe Ser
290 295 300

His Glu Asp Ser Ser Phe Gln Met Phe Ser Ser Lys Ala Tyr Ser Gln
305 310 315 320

Lys Asn Leu Leu Phe Lys Asp Ser Thr Leu Glu Leu Val Pro Ile Ala
325 330 335

Thr Gln Asn Tyr Glu Ala Trp Leu Gly Gln Glu Tyr Leu Gln Ala Met
340 345 350

Lys Gly Leu Leu Cys Asp Pro Asn Arg Leu Pro His Tyr Leu Arg Trp
355 360 365

Cys Val Leu Ser Ala Pro Glu Ile Gln Lys Cys Gly Asp Met Ala Val
370 375 380

Ala Phe Ser Arg Gln Asn Leu Lys Pro Glu Ile Gln Cys Val Ser Ala
385 390 395 400

Glu Ser Pro Glu His Cys Met Glu Gln Ile Gln Ala Gly His Thr Asp
405 410 415

Ala Val Thr Leu Arg Gly Glu Asp Ile Tyr Arg Ala Gly Lys Val Tyr
420 425 430

Gly Leu Val Pro Ala Ala Gly Glu Leu Tyr Ala Glu Glu Asp Arg Ser
435 440 445

Asn Ser Tyr Phe Val Val Ala Val Ala Arg Arg Asp Ser Ser Tyr Ser
450 455 460

Phe Thr Leu Asp Glu Leu Arg Gly Lys Arg Ser Cys His Pro Tyr Leu
465 470 475 480

Gly Ser Pro Ala Gly Trp Glu Val Pro Ile Gly Ser Leu Ile Gln Arg
485 490 495

Gly Phe Ile Arg Pro Lys Asp Cys Asp Val Leu Thr Ala Val Ser Gln
500 505 510

Phe Phe Asn Ala Ser Cys Val Pro Val Asn Asn Pro Lys Asn Tyr Pro
515 520 525

[illegible]

Sequence Listing

<110> CHUGAI SEIYAKU KABUSHIKI KAISHA
 <120> CHONDROGENESIS STIMULATOR
 5 <130> YCT-501
 <160> 15

 <210> 1
 <211> 2388
 10 <212> cDNA
 <213> Rabbit
 <400> 1
 gccgccgctc actcgttcgc actcggactc agaccagtc cgacccctg gactgcgcca 60
 tgcgggtgcg aagcgcggct atgtggatct tcctggccct gcgcaccgca ctcggcagcg 120
 15 tggaggtgcg gtggtgcacc gcgtccgagc ccgagcagca gaagtgcgag gacatgagcc 180
 aggccttccg cgaagccggc ctccagcccg cctgtctgtg cgtgcagggc acctcggccg 240
 accactgcgt ccagctcatc gcggcccacg aggcgcagc catcactctg gacggaggag 300
 ccatttacga ggcggggaag gaacacggcc tgaagcccggt ggtgggcgaa gtgtatgacc 360
 aagaggtggg cacctcctac tacgtgtgg ccgtgggtcaa gaggagctcc aacgtgacca 420
 20 tcaacaccct gagaggcgtg aagtcctgcc acacgggcat caaccgcacg gtgggctgga 480
 acgtgcctgt gggctacctg gtggacagcg gccgcctctc agtgatgggc tgtgacgtgc 540
 tcaaagcggc cagcgagtac ttcgggggca gctgcgtccc tggggcagga gagaccagat 600
 actcggagtc cctctgtcgc ctctgcggg gcgacacctc cggggagggg gtgtgtgaca 660
 agagccccct ggagcggtag tacgactaca gcggggcctt ccggtgcctg gcagaaggcg 720
 25 caggggacgt ggcctttgtg aagcacagca cggtgctgga gaacacggat gggagaacac 780
 tgccctcctg gggccacatg ctgatgtcac gggactttga gctgctgtgc cgggacggca 840
 gccgggccag cgtcaccgag tggcagcact gccacctggc ccgggtgccc gccacggcg 900
 tgggtgtccg ggccgacacc gacgcaggcc tcattctccg gcttctcaat gagggccagc 960
 ggctgttcag ccacgagggc agcagcttcc agatgttcag ctcggaggcc tacggccaga 1020

agaacctgct gttcaaagac tccacgctgg agctggtgcc catcgccaca cagacctacg 1080
 aggcctggct gggccccgag tacctgcacg ccatgaaggg tctgctctgt gacccaacc 1140
 ggctgcccc atacctgcgc tggcgcgtgc tgtccacccc cgagatccag aagtgtggag 1200
 acatggccgt ggccttcagc cggcagaggg tcaagccgga gatccagtgt gtctcggcgg 1260
 5 agtcccccca gcactgcatg gagcagatcc aggcctgggca catcgatgct gtgaccctga 1320
 acggggagga cattcacaca gcggggaaga cttatgggct gatcccggt gccggggagc 1380
 tgtatgccgc ggacgacagg agtaactcgt acttcgtggt ggccgtggtg aagcgagaca 1440
 gcgcctacgc cttaccctg gagcagctgc gcgggaagcg ctctgccac cccggcttcg 1500
 gcagcccggc cggctgggac gtcccgtgg gcgcctcat ccaactggggc tacatccggc 1560
 10 ccaggaactg cgacgtcctc acagcgggtg gtcagttctt caacgccagc tgtgtgccgg 1620
 tgaacaacc caagaagtac ccctcctcgc tgtgcgcact ctgcgtgggt gacgagcagg 1680
 gccgcaacaa gtgcaactgg aacagccagg agcggtaacta tggcgacagt ggcgcttca 1740
 ggtgcctggt ggaggggtgca ggggacgtgg ccttcgtcaa gcacacgacc atctttgaca 1800
 acacaaatgg ccacaatccc gagccgtggg ctgccatct gaggagccag gactacgagc 1860
 15 tgctgtgccc caacggcgcg cgagctgagg cgcaccagtt tgccgcctgc aacctggccc 1920
 agattccgtc ccacgccgtc atggtgcggc ccgacaccaa catcttcacc gtttacggac 1980
 tgctggacaa ggcccaggac ctgtttggag acgaccacaa caagaacggg ttcaagatgt 2040
 tcgactcctc cagctaccac ggccgagacc tgctcttcaa ggacgccag gtgcgcgctg 2100
 tgctgtggg cgagaggacc acctaccagg actggctggg gccggactac gtggcggctc 2160
 20 tggaagggat gcagtcacag cggcgtcag gggcagccgt cggcgcccc ggggcctcgc 2220
 tgctgccgt gctgccctg gctgcgggc tctgctgtc ttcgctctga gacgagcccc 2280
 gggcagcctc ggccccggca ggggagcctg cgcggaagct tctgaacga gcccgcgccc 2340
 tggctggatg tggttacctc ggcgagccgc ggggcgcgc tcccccg 2388

- 25 <210> 2
 <211> 736
 <212> PRT
 <213> Rabbit
 <400> 2

Met Arg Cys Arg Ser Ala Ala Met Trp Ile Phe Leu Ala Leu Arg Thr
1 5 10 15
Ala Leu Gly Ser Val Glu Val Arg Trp Cys Thr Ala Ser Glu Pro Glu
20 25 30
5 Gln Gln Lys Cys Glu Asp Met Ser Gln Ala Phe Arg Glu Ala Gly Leu
35 40 45
Gln Pro Ala Leu Leu Cys Val Gln Gly Thr Ser Ala Asp His Cys Val
50 55 60
Gln Leu Ile Ala Ala His Glu Ala Asp Ala Ile Thr Leu Asp Gly Gly
10 65 70 75 80
Ala Ile Tyr Glu Ala Gly Lys Glu His Gly Leu Lys Pro Val Val Gly
85 90 95
Glu Val Tyr Asp Gln Glu Val Gly Thr Ser Tyr Tyr Ala Val Ala Val
100 105 110
15 Val Lys Arg Ser Ser Asn Val Thr Ile Asn Thr Leu Arg Gly Val Lys
115 120 125
Ser Cys His Thr Gly Ile Asn Arg Thr Val Gly Trp Asn Val Pro Val
130 135 140
Gly Tyr Leu Val Asp Ser Gly Arg Leu Ser Val Met Gly Cys Asp Val
20 145 150 155 160
Leu Lys Ala Val Ser Glu Tyr Phe Gly Gly Ser Cys Val Pro Gly Ala
165 170 175
Gly Glu Thr Arg Tyr Ser Glu Ser Leu Cys Arg Leu Cys Arg Gly Asp
180 185 190
25 Thr Ser Gly Glu Gly Val Cys Asp Lys Ser Pro Leu Glu Arg Tyr Tyr
195 200 205
Asp Tyr Ser Gly Ala Phe Arg Cys Leu Ala Glu Gly Ala Gly Asp Val
210 215 220
Ala Phe Val Lys His Ser Thr Val Leu Glu Asn Thr Asp Gly Arg Thr

Phe Thr Val Asp Glu Leu Arg Gly Lys Arg Ser Cys His Pro Gly Phe
 465 470 475 480
 Gly Ser Pro Ala Gly Trp Asp Val Pro Val Gly Ala Leu Ile His Trp
 485 490 495
 5 Gly Tyr Ile Arg Pro Arg Asn Cys Asp Val Leu Thr Ala Val Gly Gln
 500 505 510
 Phe Phe Asn Ala Ser Cys Val Pro Val Asn Asn Pro Lys Lys Tyr Pro
 515 520 525
 Ser Ser Leu Cys Ala Leu Cys Val Gly Asp Glu Gln Gly Arg Asn Lys
 10 530 535 540
 Cys Thr Gly Asn Ser Gln Glu Arg Tyr Tyr Gly Asp Ser Gly Ala Phe
 545 550 555 560
 Arg Cys Leu Val Glu Gly Ala Gly Asp Val Ala Phe Val Lys His Thr
 565 570 575
 15 Thr Ile Phe Asp Asn Thr Asn Gly His Asn Pro Glu Pro Trp Ala Ala
 580 585 590
 His Leu Arg Ser Gln Asp Tyr Glu Leu Leu Cys Pro Asn Gly Ala Arg
 595 600 605
 Ala Glu Ala His Gln Phe Ala Ala Cys Asn Leu Ala Gln Ile Pro Ser
 20 610 615 620
 His Ala Val Met Val Arg Pro Asp Thr Asn Ile Phe Thr Val Tyr Gly
 625 630 635 640
 Leu Leu Asp Lys Ala Gln Asp Leu Phe Gly Asp Asp His Asn Lys Asn
 645 650 655
 25 Gly Phe Lys Met Phe Asp Ser Ser Ser Tyr His Gly Arg Asp Leu Leu
 660 665 670
 Phe Lys Asp Ala Thr Val Arg Ala Val Pro Val Gly Glu Arg Thr Thr
 675 680 685
 Tyr Gln Asp Trp Leu Gly Pro Asp Tyr Val Ala Ala Leu Glu Gly Met

gaggcgtggc tgggccatga gtacctgcac gccatgaagg gtctgctctg tgaccccaac 1140
 cggctgcccc cctacctgcg ctggtgtgtg ctctccactc ccgagatcca gaagtgtgga 1200
 gacatggccg tggccttcgc ccggcagcgc ctcaagccag agatccagtg cgtgtcagcc 1260
 aagtcccccc aacctgcat ggagcggatc caggctgagc aggtcgacgc tgtgacccta 1320
 5 agtggcgagg acatttacac ggcggggaag aagtacggcc tggttccgc agccggcgag 1380
 cactatgccc cggaagacag cagcaactcg tactacgtgg tggcgtggt gagacgggac 1440
 agtcccacg cttcacctt ggatgagctt cggggcaagc gctcctgcca cgccggtttc 1500
 ggcagccctg caggctggga tgtccccgtg ggtgccctta ttcagagagg cttcatccgg 1560
 cccaaggact gtgacgtcct cacagcagtg agcgagttct tcaatgccag ctgctgccc 1620
 10 gtgaacaacc ccaagaacta cccctcctcg ctgtgtgcac tgtgctggg ggacgagcag 1680
 ggccgcaaca agtgtgtggg caacagccag gagcggattt acggctaccg cggcgccttc 1740
 aggtgcctgg tggagaatgc gggtgacgtt gccttcgtca ggcacacaac cgtctttgac 1800
 aacacaaacg gccacaattc cgagccctgg gctgctgagc tcaggtcaga ggactatgaa 1860
 ctgctgtgcc ccaacggggc ccgagccgag gtgtcccagt ttgcagcctg caacctggca 1920
 15 cagataccac cccacgccgt gatggtccgg cccgacacca acatcttcac cgtgtatgga 1980
 ctgctggaca aggcccagga cctgtttgga gacgaccaca ataagaacgg gttcaaaatg 2040
 ttcgactcct ccaactatca tggccaagac ctgcttttca aggatgccac cgtccggggc 2100
 gtgcctgtcg gagagaaaac cacctaccgc ggctggctgg ggctggacta cgtggcggcg 2160
 ctggaaggga tgtcgtctca gcagtgtcg ggcgcagcgg ccccggcgcc cggggcgccc 2220
 20 ctgctccgc tgctgtgcc cgccctcgcc gccgcctgc tccgcccgc cctctgagcc 2280
 cggccgcccc gccccagagc tccgatgcc gcccggggag tttccgcggc ggcctctgc 2340
 gctgcggaat ccagaaggaa gctcgca 2368

<210> 4

25 <211> 738

<212> PRT

<213> Human

<400> 4

Met Arg Gly Pro Ser Gly Ala Leu Trp Leu Leu Leu Ala Leu Arg Thr

$\frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} f(x) e^{-x^2} dx = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} f(x) e^{-x^2} dx$

Leu Pro Ser Trp Gly Gln Ala Leu Leu Ser Gln Asp Phe Glu Leu Leu
 245 250 255
 Cys Arg Asp Gly Ser Arg Ala Asp Val Thr Glu Trp Arg Gln Cys His
 260 265 270
 5 Leu Ala Arg Val Pro Ala His Ala Val Val Val Arg Ala Asp Thr Asp
 275 280 285
 Gly Gly Leu Ile Phe Arg Leu Leu Asn Glu Gly Gln Arg Leu Phe Ser
 290 295 300
 His Glu Gly Ser Ser Phe Gln Met Phe Ser Ser Glu Ala Tyr Gly Gln
 10 305 310 315 320
 Lys Asp Leu Leu Phe Lys Asp Ser Thr Ser Glu Leu Val Pro Ile Ala
 325 330 335
 Thr Gln Thr Tyr Glu Ala Trp Leu Gly His Glu Tyr Leu His Ala Met
 340 345 350
 15 Lys Gly Leu Leu Cys Asp Pro Asn Arg Leu Pro Pro Tyr Leu Arg Trp
 355 360 365
 Cys Val Leu Ser Thr Pro Glu Ile Gln Lys Cys Gly Asp Met Ala Val
 370 375 380
 Ala Phe Arg Arg Gln Arg Leu Lys Pro Glu Ile Gln Cys Val Ser Ala
 20 385 390 395 400
 Lys Ser Pro Gln His Cys Met Glu Arg Ile Gln Ala Glu Gln Val Asp
 405 410 415
 Ala Val Thr Leu Ser Gly Glu Asp Ile Tyr Thr Ala Gly Lys Lys Tyr
 420 425 430
 25 Gly Leu Val Pro Ala Ala Gly Glu His Tyr Ala Pro Glu Asp Ser Ser
 435 440 445
 Asn Ser Tyr Tyr Val Val Ala Val Val Arg Arg Asp Ser Ser His Ala
 450 455 460
 Phe Thr Leu Asp Glu Leu Arg Gly Lys Arg Ser Cys His Ala Gly Phe

	465	470	475	480
	Gly Ser Pro Ala Gly Trp Asp Val Pro Val Gly Ala Leu Ile Gln Arg			
	485	490	495	
	Gly Phe Ile Arg Pro Lys Asp Cys Asp Val Leu Thr Ala Val Ser Glu			
5	500	505	510	
	Phe Phe Asn Ala Ser Cys Val Pro Val Asn Asn Pro Lys Asn Tyr Pro			
	515	520	525	
	Ser Ser Leu Cys Ala Leu Cys Val Gly Asp Glu Gln Gly Arg Asn Lys			
	530	535	540	
10	Cys Val Gly Asn Ser Gln Glu Arg Tyr Tyr Gly Tyr Arg Gly Ala Phe			
	545	550	555	560
	Arg Cys Leu Val Glu Asn Ala Gly Asp Val Ala Phe Val Arg His Thr			
	565	570	575	
	Thr Val Phe Asp Asn Thr Asn Gly His Asn Ser Glu Pro Trp Ala Ala			
15	580	585	590	
	Glu Leu Arg Ser Glu Asp Tyr Glu Leu Leu Cys Pro Asn Gly Ala Arg			
	595	600	605	
	Ala Glu Val Ser Gln Phe Ala Ala Cys Asn Leu Ala Gln Ile Pro Pro			
	610	615	620	
20	His Ala Val Met Val Arg Pro Asp Thr Asn Ile Phe Thr Val Tyr Gly			
	625	630	635	640
	Leu Leu Asp Lys Ala Gln Asp Leu Phe Gly Asp Asp His Asn Lys Asn			
	645	650	655	
	Gly Phe Lys Met Phe Asp Ser Ser Asn Tyr His Gly Gln Asp Leu Leu			
25	660	665	670	
	Phe Lys Asp Ala Thr Val Arg Ala Val Pro Val Gly Glu Lys Thr Thr			
	675	680	685	
	Tyr Arg Gly Trp Leu Gly Leu Asp Tyr Val Ala Ala Leu Glu Gly Met			
	690	695	700	

<212> DNA
 <213> Artificial Sequence
 <400> 8
 gtccggcccg acaccaacat cttc
 5
 <210> 9
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 10 <400> 9
 ggtgtgttga ggggcgtgga ctct
 <210> 10
 <211> 25
 15 <212> DNA
 <213> Artificial Sequence
 <400> 10
 tcaccaacgg ctttgagcac atcac
 20 <210> 11
 <211> 614
 <212> RNA
 <213> Mouse
 <400> 11
 25 tcaccaacgg ctttgagcac atcacagccc atcactgaca gatggccgct ctctacgagg 60
 taaccgacag gcacgttcca gccacagtc cggttaatgc ctgtgtggca ggacttgacg 120
 cccttcaggg tgttgatggt aacattggaa ttctcctga ccacagccac ggcataatag 180
 gaagtcccaa tgtcttggtc atagacttcc cccaccactg gcttcaggcc gtgctccttc 240
 cctgcctcat agatggcccc tccatccagg gtgatggcat ctgctttttg ttcttgatg 300

agctggacac agtggtcagc ggagttgccc tggacgcaga gaaggggaagg acgaatgcca 360
gctccctgga aggcctcgct catgtctttg cacttctgct gctctgcgct tgagatggta 420
caccactgca cctccatcac acagacgaca gtgcgcaggg acaggagtag ccaaaaagtc 480
acgctcagga gcctcatggc aacgttgggt tggctggggg gctggcgggg ctgtcctggc 540
5 ttctcttcc ctggtctctc tggccttcac tatttaagcg cagccccggg agagtccacg 600
ccctcaaca cacc 614

<210> 12
<211> 17
10 <212> DNA
<213> Artificial Sequence
<400> 12
tgctacttca tcgaccc

15 <210> 13
<211> 19
<212> DNA
<213> Artificial Sequence
<400> 13

20 aaagacctcc cctccatct

<210> 14
<211> 4158
<212> cDNA
25 <213> Mouse
<400> 14

ggtgtgttga ggggcgtgga ctctccccgg gctgcgctta aatagtgaag gccagagaga 60
ccaggggaaga ggaagccagg acagacccgc cagcacccca gccaacccaa cgttgccatg 120
aggctcctga gcgtgacttt ttggctactc ctgtccctgc gcactgtcgt ctgtgtgatg 180

	340	345	350
	Lys Gly Leu Leu Cys Asp Pro Asn Arg Leu Pro His Tyr Leu Arg Trp		
	355	360	365
	Cys Val Leu Ser Ala Pro Glu Ile Gln Lys Cys Gly Asp Met Ala Val		
5	370	375	380
	Ala Phe Ser Arg Gln Asn Leu Lys Pro Glu Ile Gln Cys Val Ser Ala		
	385	390	395 400
	Glu Ser Pro Glu His Cys Met Glu Gln Ile Gln Ala Gly His Thr Asp		
	405	410	415
10	Ala Val Thr Leu Arg Gly Glu Asp Ile Tyr Arg Ala Gly Lys Val Tyr		
	420	425	430
	Gly Leu Val Pro Ala Ala Gly Glu Leu Tyr Ala Glu Glu Asp Arg Ser		
	435	440	445
	Asn Ser Tyr Phe Val Val Ala Val Ala Arg Arg Asp Ser Ser Tyr Ser		
15	450	455	460
	Phe Thr Leu Asp Glu Leu Arg Gly Lys Arg Ser Cys His Pro Tyr Leu		
	465	470	475 480
	Gly Ser Pro Ala Gly Trp Glu Val Pro Ile Gly Ser Leu Ile Gln Arg		
	485	490	495
20	Gly Phe Ile Arg Pro Lys Asp Cys Asp Val Leu Thr Ala Val Ser Gln		
	500	505	510
	Phe Phe Asn Ala Ser Cys Val Pro Val Asn Asn Pro Lys Asn Tyr Pro		
	515	520	525
	Ser Ala Leu Cys Ala Leu Cys Val Gly Asp Glu Lys Gly Arg Asn Lys		
25	530	535	540
	Cys Val Gly Ser Ser Gln Glu Arg Tyr Tyr Gly Tyr Ser Gly Ala Phe		
	545	550	555 560
	Arg Cys Leu Val Glu His Ala Gly Asp Val Ala Phe Val Lys His Thr		
	565	570	575



Thr Val Phe Glu Asn Thr Asn Gly His Asn Pro Glu Pro Trp Ala Ser
580 585 590
His Leu Arg Trp Gln Asp Tyr Glu Leu Leu Cys Pro Asn Gly Ala Arg
595 600 605
5 Ala Glu Val Asp Gln Phe Gln Ala Cys Asn Leu Ala Gln Met Pro Ser
610 615 620
His Ala Val Met Val Arg Pro Asp Thr Asn Ile Phe Thr Val Tyr Gly
625 630 635 640
Leu Leu Asp Lys Ala Gln Asp Leu Phe Gly Asp Asp His Asn Lys Asn
10 645 650 655
Gly Phe Gln Met Phe Asp Ser Ser Lys Tyr His Ser Gln Asp Leu Leu
660 665 670
Phe Lys Asp Ala Thr Val Arg Ala Val Pro Val Arg Glu Lys Thr Thr
675 680 685
15 Tyr Leu Asp Trp Leu Gly Pro Asp Tyr Val Val Ala Leu Glu Gly Met
690 695 700
Leu Ser Gln Gln Cys Ser Gly Ala Gly Ala Ala Val Gln Arg Val Pro
705 710 715 720
Leu Leu Ala Leu Leu Leu Leu Thr Leu Ala Ala Gly Leu Leu Pro Arg
20 725 730 735
Val Leu

Combined Declaration for Patent Application and Power of Attorney

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

CHONDROGENESIS STIMULATOR

the specification of which (check one)

- ☐ is attached hereto;
☐ was filed in the United States under 35 U.S.C. §111 on _____, as
 U.S. Appln. No. _____*; or
☒ was/will be filed in the U.S. under 35 U.S.C. §371 by entry into the U.S. national stage of an international (PCT) application, PCT/JP00/05590 filed Aug. 21, 2000, entry requested on _____*; national stage application received U.S. Appln. No. _____*, §371/§102(e) date _____* (* if known)

and was amended on _____ (if applicable).

(include dates of amendments under PCT Art. 19 and 34 if PCT)

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; and I acknowledge the duty to disclose to the Patent and Trademark Office (PTO) all information known by me to be material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §§ 119 and 365 of any prior foreign application(s) for patent or inventor's certificate, or prior PCT application(s) designating a country other than the U.S., listed below with the "Yes" box checked and have also identified below any such application having a filing date before that of the application on which priority is claimed:

<u>232966/1999</u>	<u>Japan</u>	<u>19/8/1999</u>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day Month Year Filed)	YES	NO
_____	_____	_____	<input type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day Month Year Filed)	YES	NO

I hereby claim the benefit under 35 U.S.C. §120 of any prior U.S. non-provisional application(s) or prior PCT application(s) designating the U.S. listed below, or under §119(e) of any prior U.S. provisional applications listed below, and, insofar as the subject matter of each of the claims of this application is not disclosed in such U.S. or PCT application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the PTO all information as defined in 37 C.F.R. §1.56(a) which occurred between the filing date of the prior application and the national filing date of this application:

_____ (Application No.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)
_____ (Application No.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)
_____ (Application No.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)

As a named inventor, I hereby appoint the following registered practitioners to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

All of the practitioners associated with Customer Number 001444

Direct all correspondence to the address associated with Customer Number 001444; i.e.,
BROWDY AND NEIMARK, P.L.L.C.
 624 Ninth Street, N.W.
 Washington, D.C. 20001-5303
 (202) 628-5197

The undersigned hereby authorizes the U.S. Attorneys or Agents appointed herein to accept and follow instructions from YUASA AND HARA as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. Attorneys or Agents and the undersigned. In the event of a change of the persons from whom instructions may be taken, the U.S. Attorneys or Agents appointed herein will be so notified by the undersigned.

Page 2 of 2 Pages

Atty. Docket:

Title: CHONDROGENESIS STIMULATOR

U.S. Application filed _____, Serial No. _____

PCT Application filed August 21, 2000, Serial No. PCT/JP00/05590

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FULL NAME OF FIRST INVENTOR <u>Yukio KATO</u>		INVENTOR'S SIGNATURE <u>Yukio Kato</u>	DATE <u>Feb. 1, 2002</u>
RESIDENT <u>Hiroshima, Japan</u> <i>gp x</i>		CITIZENSHIP <u>Japanese</u>	
POST OFFICE ADDRESS <u>3-6-9-501, Ushitawaseda, Higashi-ku, Hiroshima-shi,</u> <u>Hiroshima 732-0062 Japan</u>			
FULL NAME OF SECOND JOINT INVENTOR <u>Katsumi FUJIMOTO</u>		INVENTOR'S SIGNATURE <u>Katsumi Fujimoto</u>	DATE <u>Feb. 1, 2002</u>
RESIDENT <u>Hiroshima, Japan</u> <i>gp x</i>		CITIZENSHIP <u>Japanese</u>	
POST OFFICE ADDRESS <u>1-3-11-202, Asahi, Minami-ku, Hiroshima-shi,</u> <u>Hiroshima 734-0036 Japan</u>			
FULL NAME OF THIRD JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENT		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF FOURTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENT		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF FIFTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENT		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF SIXTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENT		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF SEVENTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENT		CITIZENSHIP	
POST OFFICE ADDRESS			

ALL INVENTORS MUST REVIEW APPLICATION AND DECLARATION BEFORE SIGNING. ALL ALTERATIONS MUST BE INITIALED AND DATED BY ALL INVENTORS PRIOR TO EXECUTION. NO ALTERATIONS CAN BE MADE AFTER THE DECLARATION IS SIGNED. ALL PAGES OF DECLARATION MUST BE SEEN BY ALL INVENTORS.

INVENTOR INFORMATION

10/049957
JC13 Rec'd PCT/PTO 19 FEB 2002

Inventor One Given Name:: Yukio
Family Name:: KATO
Postal Address Line One:: 3-6-9-501, Ushitawaseda, Higashi-ku
Postal Address Line Two:: Hiroshima-shi
City:: Hiroshima
Country:: Japan
Postal or Zip Code:: 732-0062
Citizenship Country:: Japan
Inventor Two Given Name:: Katsumi
Family Name:: FUJIMOTO
Postal Address Line One:: 1-3-11-202, Asahi, Minami-ku
Postal Address Line Two:: Hiroshima-shi
City:: Hiroshima
Country:: Japan
Postal or Zip Code:: 734-0036
Citizenship Country:: Japan

CORRESPONDENCE INFORMATION

Correspondence Customer Number:: 001444
Fax One:: 202-737-3528
Electronic Mail One:: Mail@BrowdyNeimark.com

APPLICATION INFORMATION

Title Line One:: CHONDROGENESIS STIMULATOR
Total Drawing Sheets:: 6
Formal Drawings?: Yes
Docket Number:: KATO=2
Secrecy Order in Parent Appl.?: No

REPRESENTATIVE INFORMATION

Representative Customer Number:: 1444

CONTINUITY INFORMATION

This application is a:: 371 OF
> Application One:: PCT/JP00/05590
Filing Date:: 08-21-2000

PRIOR FOREIGN APPLICATIONS

Foreign Application One:: 232966/1999
Filing Date:: 08-19-1999
Country:: Japan

Priority Claimed:: Yes

PrintEFS Version 1.0.1

Source:: PrintEFS Version 1.0.1